

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:

DANILO PORRO
MICHAEL SAUER

Serial No.: 10/606,302

Filed: June 25, 2003

For: ASCORBIC ACID PRODUCTION FROM
YEAST

Confirmation No.: 4661

Group Art Unit: 1636

Examiner: Michele K. Joike

Attorney Docket: 2027.594096/RFE
(2005941)

CUSTOMER NO. 23720

REMARKS

Commissioner for Patents
P. O. Box 1450
Alexandria, VA 22313-1450

Sir:

This paper is submitted in response to the final Office Action dated October 18, 2007, for which the three-month date for response was January 18, 2008.

A request for a one-month extension of time to respond is included herewith along with the required fee. This one-month extension will bring the due date to February 18, 2008, which is within the six-month statutory period. Should such request or fee be deficient or absent, consider this paragraph such a request and authorization to withdraw the appropriate fee under 37 C.F.R. §§ 1.16 to 1.21 from Williams, Morgan & Amerson, P.C. Deposit Account No. 50-0786/2027.594096RE.

Remarks

The Examiner rejected claim 10 under 35 U.S.C. § 112, first paragraph, as allegedly lacking enablement for failure to comply with the biological material deposit requirements of 37 CFR 1.801-1.809. Applicants traverse this rejection.

Claim 10 recites (incorporating limitations from parent claims):

[A method of generating ascorbic acid, comprising a) obtaining a recombinant yeast capable of converting an ascorbic acid precursor into ascorbic acid, b) culturing the recombinant yeast in a medium comprising an ascorbic acid precursor, thereby forming ascorbic acid, and c) isolating the ascorbic acid], wherein the yeast is selected from *S. cerevisiae* strain GRF18U; *S. cerevisiae* strain W3031B; *K. lactis* strain PM6-7A; or *Z. bailii* strain ATCC 60483.

Therefore, the question is whether *S. cerevisiae* strain GRF18U; *S. cerevisiae* strain W3031B; *K. lactis* strain PM6-7A; and *Z. bailii* strain ATCC 60483 have been acceptably deposited or their deposit is not necessary. 37 CFR 1.802(b) makes clear that "[b]iological material need not be deposited, *inter alia*, if it is known and readily available to the public or can be made or isolated without undue experimentation."

S. cerevisiae strain GRF18U was deposited by coinventor Danilo Porro on July 31, 2000 with Agricultural Research Service Culture Collection (NRRL), Microbial Genomics and Bioprocessing Research Unit, National Center for Agricultural Utilization Research, 1815 North University Street, Peoria, Illinois 61604, USA. This strain was identified as deposit number NRRL Y-30320. NRRL is an International Depositary Authority under the Budapest Treaty. On August 2, 2001, Dr. Porro declared all restrictions on access to strain NRRL Y-30320 would be removed on grant of a patent on application 09/630,983, which is the parent of the present

application. Therefore, Applicant submits *S. cerevisiae* strain GRF18U has been deposited in compliance with 37 CFR 1.801-1.809.

S. cerevisiae strain W3031B is available from the American Type Culture Collection (ATCC) as deposit number ATCC 201238. The strain's availability from ATCC is demonstrated by the webpage <http://www.atcc.org/common/catalog/numSearch/numResults.cfm>, search conducted for ATCC number 201238, accessed by the undersigned on February 11, 2008, a copy of which is attached hereto. The webpage shows this strain is "known and readily available to the public." Therefore, no deposit is required (37 CFR 1.802(b)).

K. lactis strain PM6-7A is also "known and readily available to the public" as demonstrated by the many published references that cite Chen, *et al.*, Mol. Gen. Genet. **33**: 97–105 (1992) as the source of this strain. Chen, *et al.*, are coworkers of Wésolowski-Louvel, the first named author of Yeast 8, 711-719 (1992), the reference cited by the specification at p. 28, lines 3-5, as the source of this strain. References that cite Chen, *et al.*, include Tokunaga, *et al.*, Yeast **13**:699-706 (1997); Hansbro, *et al.*, Curr. Genet. **33**:46-51 (1998); Lemaire, *et al.*, Genetics **168**: 723-731 (2004); Šarinová, *et al.*, Folia Microbiol. **52**(3), 223-229 (2007); and Goffrini, FEMS Yeast Res. 7:180-187 (2007), copies of which are attached hereto. These references show that this strain was known and readily available to the public. Therefore, no deposit required. 37 CFR 1.802(b).

Z. bailii strain ATCC 60483 is available from the American Type Culture Collection (ATCC) as deposit number ATCC 60483. The strain's availability from ATCC is demonstrated by the webpage <http://www.atcc.org/common/catalog/numSearch/numResults.cfm>, search conducted for ATCC number 60483, accessed by the undersigned on February 11, 2008, a copy

of which is attached hereto. The webpage shows this strain is "known and readily available to the public." Therefore, no deposit is required. 37 CFR 1.802(b).

In conclusion, all the yeast strains recited by claim 10 are either acceptably deposited under 37 CFR 1.801-1.809 or their deposit is not necessary under 37 CFR 1.802(b). Applicants therefore request this rejection of claim 10 be withdrawn. Applicants submit all pending claims are in condition for allowance.

Respectfully submitted,

WILLIAMS, MORGAN & AMERSON, P.C.
CUSTOMER NO. 23720

February 15, 2008

/Raymund F. Eich/
Raymund F. Eich, Ph.D.
Reg. No. 42,508

10333 Richmond, Suite 1100
Houston, Texas 77042
(713) 934-4065
(713) 934-7011 (fax)

AGENT FOR APPLICANTS

Your Discoveries
Begin with Us.

Search Catalog: -- Choose Option --

[Home](#) | [Ordering Info](#) | [Quick Order](#) | [Cart](#) | [Tech Support](#) | [Contact Us](#) | [My Account](#)
| [Standards](#) | [Products](#) | [Licensing](#) | [Services](#) | [About ATCC](#) | [Log In](#)

Product Description

Before submitting an order you will be asked to read and accept the terms and conditions of ATCC's [Material Transfer Agreement](#) or, in certain cases, an MTA specified by the depositing institution.

Customers in Europe, Australia, Canada, China, Hong Kong, India, Japan, Korea, Macau, Mexico, New Zealand, Singapore, and Taiwan, R.O.C. must contact a [local distributor](#) for pricing information and to place an order for ATCC cultures and products.

Fungi, Yeast and Yeast Genetic Stock

ATCC® Number: 60483™	Order this item	Price:	\$185.00
Organism:	<i>Zygosaccharomyces bailii</i> (Lindner) Guilliermond, teleomorph; deposited as <i>Saccharomyces bailii</i> Lindner, teleomorph		
Designations:	SG 242 [MB Y-24]	Isolation:	Imported citrus concentrate, the Netherlands
Depositors:	HM Put		
Biosafety Level: 1		Shipped:	freeze-dried
Growth Conditions:	ATCC medium 323: Malt agar medium Temperature: 24.0C		
Permits/Forms:	In addition to the MTA mentioned above, other ATCC and/or regulatory permits may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please click here for information regarding the specific requirements for shipment to your location.		

Related Products

Comments:	heat-resistant [754]
Subcollection:	Yeasts
References:	754: Put HM, et al.. Heat resistance studies of yeasts; vegetative cells versus ascospores: erythromycin inhibition of sporulation in <i>Kluyveromyces</i> and <i>Saccharomyces</i> species. J. Appl. Bacteriol. 53: 73-79, 1982.

Notices and Disclaimers

ATCC products are intended for laboratory research purposes only, unless noted otherwise. They are not intended for use in humans.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this site, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.

All prices are listed in U.S. dollars and are subject to change without notice. A discount off the current list price will be applied to most cultures for nonprofit institutions in the United States. Cultures that are ordered as test tubes or flasks will carry an additional laboratory fee. Fees for permits, shipping, and handling may apply.

You may continue your ATCC Number search by typing in your search criteria below or returning to the [ATCC Search Page](#). For more information please review the [Search Help](#).

60483

[ATCC Number Search](#)[Clear Search](#)

Home Page Archive

[Home](#) | [Ordering Info](#) | [Quick Order](#) | [Support](#) | [About ATCC](#) | [Contact Us](#)
[Privacy Policy](#) | [Terms of Use](#) | [ATCC MTA](#)

© 2008 American Type Culture Collection (ATCC).
All rights reserved.



Your Discoveries
Begin with Us.

Search Catalog: -- Choose Option --

[Home](#) | [Ordering Info](#) | [Quick Order](#) | [Cart](#) | [Tech Support](#) | [Contact Us](#) | [My Account](#)
[Standards](#) | [Products](#) | [Licensing](#) | [Services](#) | [About ATCC](#) | [Log In](#)

Product Description

Before submitting an order you will be asked to read and accept the terms and conditions of ATCC's [Material Transfer Agreement](#) or, in certain cases, an MTA specified by the depositing institution.

Customers in Europe, Australia, Canada, China, Hong Kong, India, Japan, Korea, Macau, Mexico, New Zealand, Singapore, and Taiwan, R.O.C. must contact a [local distributor](#) for pricing information and to place an order for ATCC cultures and products.

Fungi, Yeast and Yeast Genetic Stock

ATCC® Number: 201238™

[Order this item](#)

Price: \$225.00

Organism: *Saccharomyces cerevisiae* Meyen ex E.C. Hansen

Designations: W303-1B

Depositors: PP Slonimski

History: ATCC<--PP Slonimski <--R. Rothstein

Biosafety Level: 1

Shipped: frozen

**Genotype/ORF/
Gene Name:** MATalpha leu2-3 leu2-112 trp1-1 ura3-1 his3-11 his3-15 ade2-1 can1-100 [25806]

**Growth
Conditions:** ATCC medium 1245: YEPD
Temperature: 25.0C

Permits/Forms: In addition to the MTA mentioned above, other ATCC and/or regulatory permits may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please [click here](#) for information regarding the specific requirements for shipment to your location.

[Related Products](#)

Applications: transformation host [25806]

Mating Type: alpha

Karyotype: Ploidy: haploid

Subcollection: Yeasts

References: 25806: Remacha M, et al. Ribosomal acidic phosphoproteins P1 and P2 are not required for cell viability but regulate the pattern of protein expression in *Saccharomyces cerevisiae*. Mol. Cell. Biol. 15: 4754-4762, 1995. PubMed: [7651393](#)

[Notices and Disclaimers](#)

ATCC products are intended for laboratory research purposes only, unless noted otherwise. They are not intended for use in humans.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this site, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.

All prices are listed in U.S. dollars and are subject to change without notice. A discount off the current list price will be applied to most cultures for nonprofit institutions in the United States. Cultures that are ordered as test tubes or flasks will carry an additional laboratory fee. Fees for permits, shipping, and handling may apply.

You may continue your ATCC Number search by typing in your search criteria below or returning to the [ATCC Search Page](#). For more information please review the [Search Help](#).

201238

[ATCC Number Search](#)

[Clear Search](#)

Home Page Archive

[Home](#) | [Ordering Info](#) | [Quick Order](#) | [Support](#) | [About ATCC](#) | [Contact Us](#)
[Privacy Policy](#) | [Terms of Use](#) | [ATCC MTA](#)

© 2008 American Type Culture Collection (ATCC).
All rights reserved.

FEMS Yeast Research

FEMS is a Federation of European Microbiological Societies, with its main mission to advance and unify microbiology knowledge. FEMS brings together 47 member societies from 36 European countries, covering over 30 000 microbiologists. Members can apply for research fellowships, visiting scientist grants, young scientist meeting grants and/or support when organizing a meeting. These benefits are restricted to members only. FEMS facilitates exchange of scientific knowledge to all microbiologists in Europe and worldwide by publishing five microbiology journals. **FEMS journals for members:** Members of FEMS Member Societies should contact customerservices@oxon.blackwellpublishing.com to arrange online access to all FEMS journals for a very attractive price.

Publications/FEMS Central Office, Keveling Buismanweg 4, 2628 CL Delft, The Netherlands. Tel.: +31 15 269 3920; fax: +31 15 269 3921; e-mail admin.journals@fems-microbiology.org

FEMS Yeast Research is published by Blackwell Publishing Ltd, 9600 Garsington Road, Oxford OX4 2DQ, UK. Tel.: +44 (0)1865 776868; fax: +44 (0)1865 714591; e-mail: femsyr@oxon.blackwellpublishing.com

Chief Editor: Teun Boekhout (Utrecht, The Netherlands)

Editors:

Andrew J. Alspaugh (Durham, NC, USA)
Barbara M. Bakker (Amsterdam, The Netherlands)
Teun Boekhout (Utrecht, The Netherlands)
Monique Bolotin-Fukuhara (Orsay Cedex, France)
Richard Calderone (Washington, DC, USA)
Terrance G. Cooper (Memphis, TN, USA)
Guenther Daum (Graz, Austria)
Ian W. Dawes (Sydney, Australia)
André Goffeau (Louvain-La Neuve, Belgium)
David S. Goldfarb (Rochester, NY, USA)
Hyun Ah Kang (Daejeon, Korea)
Cletus P. Kurtzman (Peoria, IL, USA)
Isak S. Pretorius (Adelaide, Australia)
José Ruiz-Herrera (Irapuato, Mexico)
Lubomir Tomaska (Bratislava, Slovakia)

Han H. de Winde (Delft, The Netherlands)
Ángel Domínguez (Salamanca, Spain)
Anne M. Dranginis (Jamaica, NY, USA)
Geneviève Dujardin (Gif-sur-Yvette, France)
James C. du Preez (Bloemfontein, South Africa)
Jack W. Fell (Key Biscayne, FL, USA)
Graham H. Fleet (Sydney, Australia)
Jean Marie François (Toulouse, France)
Bettina Fries (New York, USA)
Claude Gaillardin (Thierval-Grignon, France)
Joseph J. Heijnen (Delft, The Netherlands)
Bernhard Hube (Berlin, Germany)
Stephen A. James (Norwich, UK)
Lene Jespersen (Copenhagen, Denmark)
Eric A. Johnson (Madison, WI, USA)
Frans M. Klis (Amsterdam, The Netherlands)
Tatiana Kulakovskaya (Pushchino, Russia)
Marc-André Lachance (London, ON, Canada)
Stuart Levitz (Boston, MA, USA)
Ed Louis (Nottingham, UK)
Maria C. Loureiro-Dias (Lisbon, Portugal)
Ian Macreadie (Parkville, Australia)
Frank Madeo (Graz, Austria)
Leda C. Mendonça-Hagler (Rio de Janeiro, Brazil)
Francine Messenguy (Brussels, Belgium)
Tokichi Miyakawa (Hiroshima, Japan)
Silvia Moreno (Buenos Aires, Argentina)
Fritz A. Mühlschlegel (Canterbury, UK)
Gennadi I. Naumov (Moscow, Russia)
Jens Nielsen (Lyngby, Denmark)

Merja E. Penttilä (Espoo, Finland)
Jure Piškur (Lund, Sweden)
Jesús Plá (Madrid, Spain)
Danilo Porro (Milan, Italy)
Rajendra Prasad (New Delhi, India)
Jack T. Pronk (Delft, The Netherlands)
Amparo Querol (Burjassot, Spain)
Howard Riezman (Geneve, Switzerland)
José P. Sampaio (Caparica, Portugal)
Kestutis Sasnauskas (Vilnius, Lithuania)
Uwe Sauer (Zürich, Switzerland)
Johan Schnürer (Uppsala, Sweden)
Rosane F. Schwan (Lavras, Brazil)
Rafael Sentandreu (Burjassot, Spain)
Andrei A. Sibirny (Lviv, Ukraine)
Karel Sigler (Prague, Czech Republic)
David Sinclair (Boston, MA, USA)
Keshav Singh (Buffalo, NY, USA)
Matthias Sipiczki (Debrecen, Hungary)
Derek Sullivan (Dublin, Ireland)
Hana Sychrová (Prague, Czech Republic)
Joost J. Teixeira de Mattos (Amsterdam, The Netherlands)
Johan Thevelein (Leuven, Belgium)
Michael Thumm (Göttingen, Germany)
Ida J. van der Klei (Haren, The Netherlands)
Marten Veenhuis (Haren, The Netherlands)
Kevin Verstrepen (Cambridge, MA, USA)
Graeme M. Walker (Dundee, UK)
Clifford Zeyl (Winston-Salem, NC, USA)

Members of the Editorial Board:

Lennart Adler (Göteborg, Sweden)
Feng-Wu Bai (Dalian, China)
Feng-Yan Bai (Beijing, China)
Linda F. Bisson (Davis, CA, USA)
Eckhard Boles (Frankfurt am Main, Germany)
Michael Breitenbach (Salzburg, Austria)
William C. Burhans (Buffalo, NY, USA)
Peter Coote (St Andrews, Scotland, UK)
Isabel Sá Correia (Lisbon, Portugal)
Maria C. Costanzo (Stanford, CA, USA)
James M. Cregg (Claremont, CA, USA)
Heide-Marie Daniel (Louvain-la-Neuve, Belgium)
Sylvie Dequin (Montpellier, France)

Subscription rates

FEMS Yeast Research is published in 8 issues per year. Subscription prices for 2007 are: Premium Institutional: £487 (Europe), US\$900 (The Americas), £536 (Rest of World). Customers in the UK should add VAT at 6%; customers in the EU should also add VAT at 6%, or provide a VAT registration number or evidence of entitlement to exemption. Customers in Canada should add 6% GST or provide evidence of entitlement to exemption. The Premium institutional price includes online access to current content and all online back files to 1 January 1997, where available. For other pricing options or more information about online access to Blackwell Publishing journals, including access information and terms and conditions, please visit <http://www.blackwellpublishing.com/FEMSyr>.

Delivery terms and legal title: Prices include delivery of print journals to the recipient's address. Delivery terms are Delivered Duty Unpaid (DDU); the recipient is responsible for paying any import duty or taxes. Legal title passes to the customer on despatch by our distributors.

Back issues: Single issues from current and recent volumes are available at the current single issue price from Blackwell Publishing Journals. Earlier issues may be obtained from Periodicals Service Company, 11 Main Street, Germantown, NY 12526, USA. Tel.: +1 518 537 4700; fax: +1 518 537 5899; e-mail: psc@periodicals.com

Periodical ID statement: FEMS MICROBIOLOGY ECOLOGY (ISSN: 0168-6496) is published in 12 issues per year. US mailing agent: G3 Worldwide (US) Inc., 8701 Bellanca Ave., Los Angeles, CA 90045. Periodical postage paid at Los Angeles, CA. POSTMASTER: Send all address changes to FEMS MICROBIOLOGY ECOLOGY, Blackwell Publishing Inc., Journals Subscription Department, 350 Main St., Malden, MA 02148-5020, USA.

Journal Customer Services: For ordering information, claims and any enquiry concerning your journal subscription please contact your nearest office:

UK: Email: customerservices@oxon.blackwellpublishing.com; tel: +44 (0) 1865 778315; fax: +44 (0) 1865 471775

USA: E-mail: subscrip@bos.blackwellpublishing.com; tel: +1 781 388 8206 or +1 800 835 6770 (Toll free in the USA); fax: +1 781 388 8232

Asia: E-mail: subs@blackwellpublishingasia.com; tel: +61 3 8359 1100; fax: +61 3 8359 1120

Production Editors: Marion Laws, Julie Smith and Helen Birchall (email: femsyr@oxon.blackwellpublishing.com)

Copyright and Photocopying

© 2007 Federation of European Microbiological Societies/ Blackwell Publishing Ltd. All rights reserved. No part of this publication (apart from articles marked OnlineOpen) may be reproduced, stored or

transmitted in any form or by any means without the prior permission in writing from the copyright holder. Authorization to photocopy items for internal and personal use is granted by the copyright holder for libraries and other users registered with their local Reproduction Rights Organisation (RRO), e.g. Copyright Clearance Center (CCC), 222 Rosewood Drive, Danvers, MA 01923, USA (www.copyright.com), provided the appropriate fee is paid directly to the RRO. This consent does not extend to other kinds of copying such as copying for general distribution, for advertising or promotional purposes, for creating new collective works or for resale. OnlineOpen articles are made available in accordance with the terms of the Creative Commons Deed, Attribution 2.5 (further details available from <http://www.creativecommons.org>), which allows Open Access dissemination of the article, but does not permit commercial exploitation or the creation of derivative works. Special requests should be addressed to Blackwell Publishing at: journalsrights@oxon.blackwellpublishing.com

Disclaimer

The Publisher, FEMS and Editors cannot be held responsible for errors or any consequences arising from the use of information contained in this journal; the views and opinions expressed do not necessarily reflect those of the Publisher, FEMS and Editors, neither does the publication of advertisements constitute any endorsement by the Publisher, FEMS and Editors of the products advertised.

Paper

Blackwell Publishing's policy is to use permanent paper from mills that operate a sustainable forestry policy, and which has been manufactured from pulp that is processed using acid-free and elementary chlorine-free practices. Furthermore, Blackwell Publishing ensures that the text paper and cover board used in all our journals has met acceptable environmental accreditation standards.

CarbonNeutral statement: Blackwell Publishing is a CarbonNeutral company. For more information visit <http://www.blackwellpublishing.com/carbonneutral>

Free access

Access to this journal is available free online within institutions in the developing world through the AGORA initiative with the FAO and the HINARI initiative with the WHO. For information, visit <http://www.aginternetwork.org>, <http://www.healthinternetwork.org>

Printed in Singapore by HO Printing Pte Ltd

ISSN 1567-1356 (Print)

ISSN 1567-1364 (Online)

For submission instructions, subscription and all other information visit: <http://www.blackwellpublishing.com/FEMSyr>

A respiratory-deficient mutation associated with high salt sensitivity in *Kluyveromyces lactis*

Paola Goffrini

Department of Genetics, Biology of Microorganisms, Anthropology, Evolution, University of Parma, Parco Area delle Scienze, Parma, Italy

Correspondence: Paola Goffrini,
Department of Genetics, Biology of
Microorganisms, Anthropology, Evolution,
University of Parma, Parco Area delle Scienze
11A, 43100 Parma, Italy. Tel.: +39 0521
905602; fax: +39 0521 905604; e-mail:
paola.goffrini@unipr.it

Received 24 February 2006; revised 13 June
2006; accepted 18 June 2006.
First published online 7 August 2006.

DOI:10.1111/j.1567-1364.2006.00148.x

Editor: Monique Bolotin-Fukuhara

Keywords

salt resistance; respiratory deficiency;
Kluyveromyces lactis; yeast; *BCS1*; *VMA13*.

Abstract

A salt-sensitive mutant of *Kluyveromyces lactis* was isolated that was unable to grow in high-salt media. This mutant was also respiratory-deficient and temperature-sensitive for growth. The mutation mapped in a single nuclear gene that is the ortholog of *BCS1* of *Saccharomyces cerevisiae*. The *BCS1* product is a mitochondrial protein required for the assembly of respiratory complex III. The *bcs1* mutation of *S. cerevisiae* leads to a loss of respiration, but, unlike in *K. lactis*, it is not accompanied by salt sensitivity. All the respiratory-deficient *K. lactis* mutants tested were found to be salt-sensitive compared to their isogenic wild-type strains. In the presence of the respiratory inhibitor antimycin A, the wild-type strain also became salt-sensitive. By contrast, none of the *S. cerevisiae* respiratory-deficient mutants tested showed increased salt sensitivity. The salt sensitivity of the *Klbcs1* mutant, but not its respiratory deficiency, was suppressed by the multicopy *KIVMA13* gene, a homolog of the *S. cerevisiae* *VMA13* gene encoding a subunit of the vacuolar H^+ -ATPase. These results suggest that cellular salt homeostasis in *K. lactis* is strongly dependent on mitochondrial respiratory activity, and/or that the ion homeostasis of mitochondria themselves could be a primary target of salt stress.

Introduction

Yeast species are resistant, to various degrees, to high-salt environments. In the mechanisms of this resistance, we may distinguish two processes: (1) rapid primary response of the cell to dehydration; and (2) slow adaptation to the high-salt environments.

The first process, extensively studied in *Saccharomyces cerevisiae*, appears to be associated with increased turnover/accumulation of glycerol (or polyalcohol) and trehalose (Hohmann, 2002), and with the exclusion from the cell of a harmful solute such as NaCl, mediated by the P-type ATPase Ena1p and the sodium-proton antiporter Nha1p (Serrano *et al.*, 1997; Patterson *et al.*, 1999). The second process, which involves slow adaptation to high salt, is still poorly explored. In some cases, genomic rearrangements, such as gene amplification, may be involved (Prior *et al.*, 1996; Albrecht *et al.*, 2000). Studies of salt-sensitive mutations have revealed a wide range of associated phenotypes. In *S. cerevisiae*, the expression of at least 18 genes was strongly induced by high salt (Blomberg, 1995).

In the present work, to enable the study of salt sensitivity in *Kluyveromyces lactis* we selected mutants with high salt sensitivity. Among them, we found a new kind of mutant

that showed, in addition to its sensitivity to high concentrations of NaCl, KCl and LiCl, a complete loss of respiratory activity. The mutation mapped in the *KIBCS1* gene, which codes for a mitochondrial protein involved in the assembly of respiratory complex III. A direct association of the salt sensitivity of yeast with a specific mitochondrial dysfunction has never been described previously. In the present work, the properties of the *Klbcs1* mutation and the association between salt sensitivity and mitochondrial dysfunction will be described.

Materials and methods

Strains, media and growth conditions

Table 1 lists the *K. lactis* and *S. cerevisiae* strains and derived mutants used in this study. *Escherichia coli* DH10B was used as a cloning host and for DNA propagation.

Complete medium (YP) contained 1% Bacto yeast extract (Difco) and 1% Bacto peptone (Difco). It was supplemented with a carbon source at 2% (glucose, glycerol or others as specified). Minimal medium contained 0.7% Yeast Nitrogen Base without amino acids (Difco), and 2% glucose, supplemented with appropriate auxotrophic requirements. For

Table 1. List of yeast strains

Strain	Genotype	Source
<i>K. lactis</i>		
PM6-7A	<i>MATa uraA1-1 adeT-600</i>	Chen <i>et al.</i> (1992)
PM6-7A/A16	<i>MATa uraA1-1 adeT-600 KlbcS1</i>	This study
PM6-7A/ Δ KlcyC1 (KG6)	<i>MATa uraA1-1 adeT-600, KlcyC1::URA3</i>	Chen & Clark-Walker (1993)
MW179-1D	<i>MATa uraA1-1 leu2 lac4-8 trpA1 ade1</i>	M. Wésolowski-Louvel (University of Lyon 1)
MW179-1D/ Δ Klcox14	<i>MATa uraA1-1 leu2 lac4-8 trpA1 ade1 Klcox14::kanMX4</i>	Fiori <i>et al.</i> (2000)
JBD100	<i>MATa trp1 lac4-1 ura3</i>	Heus <i>et al.</i> (1990)
JBD100/Klcox18 (M5)	<i>MATa trp1 lac4-1 ura3 Klcox18</i>	Hikkel <i>et al.</i> (1997)
JBD100/Klcytc1 (M3)	<i>MATa trp1 lac4-1 ura3 Klcytc1</i>	Gbelská <i>et al.</i> (1996)
WMH9802/ Δ Klqcr8	<i>MATa uraA1-1 leu2 lac4-8 trpA1 ade1 Klqcr8::URA3</i>	Brons <i>et al.</i> (2001)
2360/7	<i>MATa lysA</i>	Parma collection
<i>S. cerevisiae</i>		
W303-1A	<i>MATa SUC2 ade2 can1 his3 leu2 trp1 ura3</i>	R. Rothstein (Columbia University)
W303-1A/ Δ bcs1	<i>MATa SUC2 ade2 can1 his3 leu2 trp1 ura3 bcs1::HIS3</i>	Nobrega <i>et al.</i> (1992)
BY4741	<i>MATa ura3Δ0 his3Δ1 leu2Δ0 met15Δ0</i>	Euroscarf collection
BY4741/ Δ sop1 (YPR032W)	<i>MATa ura3Δ0 his3Δ1 leu2Δ0 met15Δ0 sop1::kanMX4</i>	Euroscarf collection
MH41-7B <i>rho</i> ⁺ and <i>rho</i> ⁻ derivatives*	<i>MATa ade2 his1</i>	Institut Curie, Orsay
MH32-12D <i>rho</i> ⁺ and <i>rho</i> ⁰ derivatives*	<i>MATa ade2 his1</i>	Institut Curie, Orsay
IL8-8C/HF71/ <i>rho</i> ⁰	<i>MATa trp1 his1, rho</i> ⁰	Institut Curie, Orsay
IL125-10C/ <i>rho</i> ⁰	<i>MATa ura, rho</i> ⁰	Institut Curie, Orsay

*MH41-7B/HF21 *rho*⁰, MH41-7B/OI-3 *rho*⁻, MH41-7B/C7 *rho*⁻, MH41-7B/P1 *rho*⁻, MH32-12D/*rho*⁰ (Wésolowski-Louvel & Fukuhara, 1979)

plate tests, these media were solidified with 2% agar. Salt resistance/sensitivity was tested on YP-glucose plates containing the indicated concentrations of salts or sugars. The culture temperature was 28 °C unless specified otherwise. Antimycin A (Sigma) was used at a concentration of 5 μ M throughout. Genetic procedures for mating and sporulation were done on ME plates (5% malt extract, 3% Bacto agar).

Isolation of mutants

Yeast cells were mutagenized with UV irradiation according to Wésolowski-Louvel *et al.* (1992). Cells at a density of 10⁸ cells mL⁻¹ were exposed to 75 J m⁻² of UV radiation. Survival was 20–30%. Cells were plated for single colonies on YP-glucose, and replica-plated on NaCl-containing medium. Putative mutants (negative growth on 1.5 M NaCl) were subcloned and retested for their salt-sensitive phenotype.

Cytochrome absorption spectra

Cells, grown to early stationary phase on YP medium supplemented with 2% glucose, were harvested by centrifugation, washed twice with cold (4 °C) distilled water, and suspended in a volume of cold water twice the pellet volume. Differential spectra between reduced and oxidized cells were recorded at room temperature using a Cary 219 spectrometer. The bandwidth was 1 nm and the scan speed was 0.5 nm s⁻¹. The cell suspension was reduced by sodium dithionite.

General methods

Published procedures were used for the transformation of *K. lactis* (Bianchi *et al.*, 1987) and *E. coli* (Mandel & Higa, 1970). DNA manipulation, restriction enzyme digestion, plasmid engineering and standard techniques were performed according to Sambrook & Russel (2001). Sequencing was performed using a Beckman CEQ2000 automatic sequencer. Sequence analysis was performed with the BLASTP program (Altschul *et al.*, 1990), and sequence alignment with the CLUSTAL W program (Thompson *et al.*, 1994). The GenBank accession numbers for *KIBCS1* and for *KIVMA13* are [AJ299738](#) and [AJ547613](#), respectively.

The amplification of the mutated *KlbcS1* allele was obtained by PCR with PM6-7A/A16 genomic DNA as a template and the primer pair *KIBCS1*F, 5'-AATCCGAGGCGCTCGATTTC-3', and *KIBCS1*R, 5'-GGATGGACAACGAACGATAT-3'.

Results

Isolation and phenotypic characterization of mutants of *K. lactis* with high salt sensitivity

The strain PM6-7A was UV-mutagenized, and about 25 000 cells were plated on YP-glucose plates. The colonies were replica-plated onto YP-glucose containing 1.5 M NaCl. Many colonies showed slow, leaky growth. Only three colonies (A8, A10 and A16) were clearly incapable of growing on the high-salt plates, as shown in Fig. 1. They




Fig. 1. Salt sensitivity of *Kluyveromyces lactis* mutants. Strain PM6-7A and the three salt-sensitive mutants derived from it were streaked on YP-glucose plates with (b) or without (a) 1.5 M NaCl, and allowed to grow at 28 °C for 2 days.

Table 2. Salt sensitivity of three mutants

Strain	NaCl 1.5 M	KCl 1.5 M	LiCl 0.3 M	Glucose 2 M	Sorbitol 2 M	Sucrose 1 M
PM6-7A	+	+	+	+	+	+
PM6-7A/A8	—	+	+ / —	+	+	+
PM6-7A/A10	—	+	—	+	+	+
PM6-7A/A16	—	—	—	+	+	+

The three mutants, A8, A10 and A16, are subclones of PM6-7A obtained by UV mutagenesis. All the tests were performed on YP-glucose plates supplemented with high-osmolality solutes as indicated. The signs + and — indicate the occurrence and absence of growth, respectively, as recorded after 3 days of incubation.

were submitted to further tests of osmosensitivity on high-salt and high-sugar media. The results are shown in Table 2.

All the mutants were sensitive to 1.5 M NaCl, but not to high concentrations of sugars. They could be distinguished by their different sensitivities to 1.5 M KCl and to 300 mM LiCl. In particular, a mutant, called A16, was sensitive to all three salts, and moreover its growth was sensitive to high temperature (36 °C) on YP-glucose. This mutant was also unable to grow on nonfermentable substrates (glycerol, ethanol and lactate). It was indeed respiratory-deficient, displaying an 80-fold decrease in oxygen consumption rate compared to the wild-type parental strain (data not shown). The cytochrome absorption spectra (Fig. 2) indicated that cytochrome *b* and cytochrome *a+a₃* were reduced by 25% and 65%, respectively. Our study focused on the particular pleiotropic mutant A16.

The A16 mutant appears to have a single gene mutation, as suggested by two observations: (1) all spontaneous back mutations restored a complete wild-type phenotype, and (2) transformation of the mutant with a single gene (*KIBCS1*), as described below, fully complemented all the deficient phenotypes of A16. Although genetic crosses with wild-type laboratory strains gave diploids severely impaired in sporulation, we were able to obtain a diploid able to sporulate by crossing the mutant with the 2360/7 strain. Tetrad analysis demonstrated that the pleiotropic phenotype was due to a single nuclear mutation, as we obtained a 2 : 2 Mendelian segregation.

Cloning and characterization of the *KIBCS1* gene

The A16 mutant was transformed with a *K. lactis* genomic library constructed on a centromere-based vector KCp491





Fig. 2. Cytochrome spectra of *Kluyveromyces lactis* PM6-7A and its respiratory-deficient mutant A16. The absorbance peaks at 550, 560 and 602 nm correspond to cytochrome *c*, cytochrome *b* and cytochromes *a+a₃*, respectively.

carrying a *URA3* marker (Prior *et al.*, 1993). The *Ura*⁺ transformants were tested for their salt sensitivity. Among them, two clones were capable of growing on 1.5 M NaCl. The *Ura*⁺ phenotype and the salt resistance cosegregated in the course of spontaneous loss of the marker. The two transformants also recovered temperature resistance and respiratory competence. Both carried an identical plasmid (named pOSME) that contained a DNA insert of 6.0 kb. Fragments of this DNA were subcloned, and a segment spanning a 2.7-kb *Eco*RI fragment (carried by a plasmid named pOSME/E27) was found to be responsible for the transformed phenotype. The insert contained a putative single ORF of 1.35 kb. The predicted product of this DNA

Fig. 3. Structure of KIBCS1p as deduced from the DNA sequence. The amino acid sequence of the KIBCS1 product was deduced from the nucleotide sequence. Comparison with the *Saccharomyces cerevisiae* ortholog (Nobrega *et al.*, 1992) clearly identified characteristic sequence motifs as indicated. The lower half of the figure shows the DNA sequence-deduced structure of KIBCS1p in the A16 mutant.



was a 450-amino-acid protein that showed 69% identity with the Bcs1p of *S. cerevisiae*. This is a nuclear-encoded mitochondrial protein of the AAA family (ATPase associated with diverse cellular activities) that in *S. cerevisiae* controls the assembly of the cytochrome *bc*₁ complex and leads to a total absence of complex III activity. It has been proposed that Bcs1p acts as an ATP-dependent chaperone maintaining the precomplex in a competent state for the subsequent assembly of Rieske FeS and Qcr10p proteins (Nobrega *et al.*, 1992; Cruciat *et al.*, 1999).

The complex III deficiency in the *bcs1* mutant also affects the amount of complex IV assembly and activity. In the cytochrome absorption spectra of the *Klbcs1* mutant (Fig. 2), the effect on cytochromes *a*+*a*₃ of complex IV was more evident; the effect on cytochrome *b* was less pronounced but distinguishable from that of the wild type. With regard to cytochrome *c*₁, the negative effect of the mutation is evident from the reduced shoulder on the right side of the peak at 550 nm. Taken together, these results indicated that there was the same pleiotropic effect on cytochrome assembly in the *K. lactis bcs1* mutant as was observed in the *S. cerevisiae* mutant.

As expected, the structural organization of the cloned *K. lactis* gene was similar to that of *BCS1*, showing the presence of two supposed ATP-binding motifs and a mitochondria-targeting signal (Fig. 3).

The mutant allele of the A16 strain was also sequenced by means of triplicate PCR amplification of the mutant DNA. The mutation corresponded to the introduction of a stop codon in the middle of the gene, resulting in a large deletion of the C-terminal half of the protein, with a loss of the putative ATP-binding motifs (Fig. 3).

Functional complementation of *S. cerevisiae bcs1* mutation by the cloned *K. lactis* gene

Given the structural similarity of the cloned gene to *BCS1*, we carried out a functional complementation experiment. The cloned *K. lactis* DNA on the KCp491 vector that can also

replicate in *S. cerevisiae* was transformed into the *S. cerevisiae Δbcs1* mutant. The transformants fully recovered the ability to grow on nonfermentable carbon sources (Table 3). The *K. lactis* gene was therefore named *KIBCS1*, and the mutant allele *Klbcs1*. Having observed a functional homology between *KIBCS1* and *BCS1*, we wanted to know whether in *S. cerevisiae* the *BCS1* gene is involved in salt resistance. In contrast to what was observed in *K. lactis*, the *S. cerevisiae Δbcs1* mutant did not show increased sensitivity to 1 M NaCl compared to its isogenic wild type (1 M was the highest concentration tolerated by the wild-type *S. cerevisiae* strains used here, as compared to 1.5 M for *K. lactis* strains) (Table 3). Moreover, the mutant was also not temperature-sensitive for growth. Thus the phenotypes of the *bcs1* mutation clearly differed between the two species.

Does the salt-sensitive phenotype always accompany the respiratory deficiency in *K. lactis*?

In order to know whether the salt sensitivity was due to the *Klbcs1* mutation *per se* or to the respiratory deficiency resulting from the mutation, we examined the salt sensitivity of available *K. lactis* respiratory-deficient mutants. Several respiratory-deficient mutants of *S. cerevisiae* were also included for comparison. The results obtained (Table 3) indicated that all the respiratory-deficient mutants of *K. lactis* were sensitive to 1.5 M NaCl (Fig. 4). These were mutants of cytochrome *c*, cytochrome oxidase subunits and complex III subunits, respectively. One exception was a cytochrome *c*₁ mutant that grew on the high-salt medium (see Discussion). The strong correlation between respiratory deficiency and salt sensitivity was further supported by the observation that the wild-type *K. lactis* strains became salt-sensitive in the presence of the respiratory inhibitor antimycin A (Table 3). By contrast, none of the respiratory-deficient mutants of *S. cerevisiae*, including *bcs1*, were salt-sensitive compared to isogenic wild-type strains. As a

Table 3. Respiratory deficiency and salt sensitivity: comparison between *Kluyveromyces lactis* and *Saccharomyces cerevisiae*

	YP-glycerol	YP-glucose + NaCl
<i>K. lactis</i>		
PM6-7A	+	+
PM6-7A/A16	—	—
PM6-7A/A16+[<i>KIBC51</i>]	+	+
PM6-7A/ Δ Klcy1	—	—
MW179-1D	+	+
MW179-1D/ Δ Klcox14	—	—
WMH9802/ Δ Klqcr8	—	—
JBD100	+	+
JBD100/Klcox18	—	—
JBD100/Klcytc1	—	+
PM6-7A+antimycin A	—	—
MW179-1D+antimycin A	—	—
JBD100+antimycin A	—	—
<i>S. cerevisiae</i>		
W303-1A	+	+
W303-1A/ Δ bcs1	—	+
W303-1A/ Δ bcs1+[<i>KIBC51</i>]	+	+
MH41-7B, ρ^+	+	+
MH41-7B/HF21, ρ^0	—	+
MH41-7B/OI-3, ρ^-	—	+
MH41-7B/C7, ρ^-	—	+
MH41-7B/P1, ρ^-	—	+
MH32-12D, ρ^+	+	+
MH32-12D/ ρ^0	—	+
IL8-8C/HF71, ρ^0	—	+
IL125-10C/ ρ^0	—	+
BY4741	+	+
BY4741/ Δ sop1	+	—
W303-1A+antimycin A	—	+
BY4741+antimycin A	—	+

Kluyveromyces lactis and *Saccharomyces cerevisiae* strains were streaked on YP plates containing glycerol (test for respiratory competence) or glucose and NaCl (test for salt sensitivity), as indicated. +[*KIBC51*] indicates the presence of the monocopy plasmid carrying the *KIBC51* gene. Note that for *S. cerevisiae*, the salt sensitivity test was performed on 1 M NaCl, the maximal salt concentration tolerated by the wild-type strains, as compared to 1.5 M for *K. lactis*. The *K. lactis* wild-type strains used here cease to grow on 1.7 M NaCl.

negative control, we used the *sop1* mutant of *S. cerevisiae*, a well-known salt-sensitive strain (Larsson *et al.*, 1998).

A multicopy suppressor of *Klbc51*

The role of KlBcs1p in salt resistance is not obvious. In order to find possible linked elements, we looked for a multicopy suppressor of *Klbc51* mutation. The *Klbc51* mutant was transformed with a *K. lactis* genomic library carried by the multicopy *K. lactis*/*S. cerevisiae* shuttle vector pSK1 (Wésolowski-Louvel *et al.*, 1988). Among the 6000 Ura⁺ transformants, one single clone recovered the ability to grow on 1.5 M NaCl (in YP-glucose). However, this clone re-

mained respiratory-deficient and temperature-sensitive for growth. Therefore, the suppressor appeared to be extragenic. The suppressed clone contained a plasmid with a DNA insert of about 6 kb. The predicted product of the gene found in this segment has an identity of 43% with the protein encoded by the *VMA13* gene of *S. cerevisiae*. The Vma13p of *S. cerevisiae* has been known to form part of the vacuolar H⁺-ATPase complex (V-ATPase), an ATP-dependent proton pump that acidifies the vacuolar compartment. Vma13p is thought to be an activator or a stabilizer of the multimeric V-ATPase complex (Anraku *et al.*, 1992). The functional equivalence of *KIVMA13* and *VMA13* was confirmed by complementation of the *vma13* mutation by *KIVMA13* (recovery from Ca²⁺ sensitivity of the mutant; data not shown).

A contribution of V-ATPase to the mechanism of salt tolerance in yeast has been reported (Hamilton *et al.*, 2002). ATP hydrolysis is coupled with active proton transport inside the vacuole, thus generating a chemical gradient that drives transport of ions such as Na⁺ and Ca²⁺, which is mediated by the Na⁺/H⁺ antiporter, NHX1 (Nass & Rao, 1998; Hirata *et al.*, 2002).

If the observed suppression resulted from a major role played by V-ATPase in the ionic homeostasis of mitochondria, we might expect that the salt-sensitive phenotype of other *K. lactis* respiratory-deficient mutants might also be complemented by the multicopy *KIVMA13* gene. However, this was not the case. The suppressor effect of *KIVMA13* was specific for the *Klbc51* mutation, suggesting that if the V-ATPase contributes to the ionic balance of the mitochondria, this might occur through a mechanism involving Bcs1p.

Discussion

Relationship between respiratory deficiency and salt resistance in *K. lactis*

Salt resistance involves many genes. Studies on several yeast species, in particular *S. cerevisiae*, have led to the identification of different types of mutation that display different phenotypes. A striking finding was the identification of *BCS1* as a genetic determinant involved in salt resistance in *K. lactis*. Because it is difficult to imagine a specific direct involvement of Bcs1p in salt resistance, we hypothesized that the salt sensitivity of the *Klbc51* mutant may not be a specific phenotype of this particular mutation, but rather a general consequence of the respiratory deficiency. The observation that wild-type *K. lactis* strains became salt-sensitive when respiration was specifically blocked by antimycin A was in favor of this interpretation. When we investigated the osmotic response of other respiratory-deficient mutants of *K. lactis*, all of them, as expected, were salt-sensitive, except




Fig. 4. Salt-sensitive phenotype of *Kluyveromyces lactis* respiratory-deficient mutants. The salt sensitivity of various respiratory mutants of *K. lactis* was tested against either 1.5 M NaCl or 1.5 M KCl included on YP-glucose plates. The respiratory deficiency is shown by their absence of growth on YP-glycerol plates.

for a cytochrome c_1 mutant that maintained the resistant phenotype of the parental strain. However, this mutant retained 20% of the respiratory capacity of its wild-type parent (Gbelská *et al.*, 1996). Thus, this mutant may not be considered to be strictly respiration-negative. For this reason, it should not be considered an exception to the relationship between respiratory deficiency and salt sensitivity. Therefore, the results with all the respiratory-deficient mutants, as well as those obtained with antimycin A, indicate that the lack of respiratory activity was correlated with the salt sensitivity.

Difference between *K. lactis* and *S. cerevisiae*

Unlike in *K. lactis*, in *S. cerevisiae* the *bcs1* mutant as well as all other respiratory-deficient mutants retain high salt resistance, which is similar to the level observed in the wild-type strains. Therefore, the two yeast species clearly differ by the presence/absence of a link between respiratory deficiency and salt sensitivity. A possible reason could be that the laboratory strains of *S. cerevisiae* originate mainly from fermentation media, and hence these strains show a preference for a fermentative life rather than a respiratory

mode of growth. We would expect such yeasts to possess stress-resisting mechanisms that do not require respiratory metabolism. Conversely, *K. lactis* strains have a strong respiratory activity, and normally this microorganism has a respiratory mode of life. The reducing potential generated by its strong glucose 6-phosphate shunt has to be recycled by active respiratory activity. Therefore, this species may have developed a stress response mechanism that is more tightly associated with mitochondrial functions, in comparison with *S. cerevisiae*. A respiratory deficiency or a mitochondrial mutation may then result in increased sensitivity to certain stresses. Such an interpretation, perhaps oversimplified, can be experimentally tested by the use of other yeast species showing a high dependence of growth on respiratory activity. In those species, salt resistance may also be linked to active mitochondrial functions, and in this regard, the responses of *S. cerevisiae* to high salt may reflect an exceptional physiology of this species.

Finally, the specific suppression of multicopy *KIVMA13* in the *Klbcsl* mutation has not yet been explained. Moreover, a relationship of V-ATPase with the mitochondrial system has been reported by Ohya *et al.* (1991), who observed that some of the mutations of the V-ATPase complex were accompanied

by a respiratory-deficient phenotype in *S. cerevisiae*. Recently, links between iron and copper metabolism and mitochondrial and vacuolar function have also been found. In particular, a role for *VMA13* in metal trafficking has been demonstrated (van Bakel *et al.*, 2005).

Acknowledgements

I would like to thank Hiroshi Fukuhara for encouragement and helpful discussions, and Marco Ventura for critical reading of the manuscript. I thank Drs Julius Šubík, Alexander Tzagoloff, Xin-Jie Chen, Janike Brons, Claudio Falcone and Micheline Wésolowski-Louvel for providing mutant strains. I thank Tiziana Monduzzi for skillful technical assistance. This work was supported by Fondazione Adriano Buzzati-Traverso.

References

- Albrecht EB, Hunyady AB, Stark GR & Patterson TE (2000) Mechanisms of *sod2* gene amplification in *Schizosaccharomyces pombe*. *Mol Biol Cell* 3: 873–886.
- Altshul SF, Gish W, Miller W, Myers EW & Lipman DJ (1990) Basic local alignment search tool. *J Mol Biol* 215: 403–410.
- Anraku Y, Hirata R, Wada Y & Ohya Y (1992) Molecular genetics of the yeast vacuolar H⁺-ATPase. *J Exp Biol* 172: 67–81.
- Bianchi MM, Falcone C, Chen XJ, Wésolowski-Louvel M, Frontali L & Fukuhara H (1987) Transformation of the yeast *Kluyveromyces lactis* by new vectors derived from the 1.6 µm circular plasmid pKD1. *Curr Genet* 12: 185–192.
- Blomberg A (1995) Global changes in protein synthesis during adaptation to 0.7 M NaCl medium of *Saccharomyces cerevisiae*. *J Bacteriol* 177: 3563–3572.
- Brons JF, Dryla AA, Pluger EB, Vinkenvleugel TM, Hornig NC, Grivell LA & Blom J (2001) Carbon source-dependent transcriptional regulation of the *QCR8* gene in *Kluyveromyces lactis*: identification of *cis*-acting regions and *trans*-acting factors in the *KIQCR8* upstream region. *Curr Genet* 39: 311–318.
- Chen XJ & Clark-Walker GD (1993) Mutations in *MGI* genes convert *Kluyveromyces lactis* into a petite-positive yeast. *Genetics* 133: 517–525.
- Chen XJ, Wésolowski-Louvel M & Fukuhara H (1992) Glucose transport in the yeast *Kluyveromyces lactis*. II. Transcriptional regulation of the glucose transporter gene *RAG1*. *Mol Gen Genet* 33: 97–105.
- Cruciat CM, Hell K, Fölsch H, Neupert W & Stuart RA (1999) Bcs1p, an AAA-family member, is a chaperone for the assembly of the cytochrome *bc1* complex. *EMBO J* 18: 5226–5233.
- Fiori A, Saliola M, Goffrini P & Falcone C (2000) Isolation and molecular characterization of *KICOX14*, a gene of *Kluyveromyces lactis* encoding a protein necessary for the assembly of the cytochrome oxidase complex. *Yeast* 16: 307–314.
- Gbelská Y, Horváthová K, van der Aart QJ, Zonneveld BJ, Steensma HY & Šubík J (1996) Isolation and molecular analysis of the gene for cytochrome *c₁* of *Kluyveromyces lactis*. *Curr Genet* 30: 145–150.
- Hamilton CA, Taylor GJ & Good AG (2002) Vacuolar H⁺-ATPase, but not mitochondrial F₁F₀-ATPase, is required for NaCl tolerance in *Saccharomyces cerevisiae*. *FEMS Microbiol Lett* 208: 227–232.
- Heus JJ, Zonneveld BJ, Steensma HY & van den Berg JA (1990) Centromeric DNA of *Kluyveromyces lactis*. *Curr Genet* 18: 517–522.
- Hikkel I, Gbelská Y, van der Aart QJ, Lubecu G & Šubík J (1997) Cloning and characterization of *KICOX18*, a gene required for activity of cytochrome oxidase in *Kluyveromyces lactis*. *Curr Genet* 32: 267–272.
- Hirata T, Wada Y & Futai M (2002) Sodium and sulfate ion transport in yeast vacuoles. *J Biochem* 131: 261–265.
- Hohmann S (2002) Osmotic stress signaling and osmoadaptation in yeasts. *Microbiol Mol Biol Rev* 66: 300–372.
- Larsson K, Böhl F, Sjöström I, Akhtar N, Strand D, Mechler BM, Grabowski R & Adler L (1998) The *Saccharomyces cerevisiae* *SOP1* and *SOP2* genes, which act in cation homeostasis, can be functionally substituted by the *Drosophila lethal(2)giant larvae* tumor suppressor gene. *J Biol Chem* 273: 33610–33618.
- Mandel JK & Higa S (1970) Calcium dependent bacteriophage DNA infection. *J Mol Biol* 53: 159–162.
- Nass R & Rao R (1998) Novel localization of a Na⁺/H⁺ exchanger in a late endosomal compartment of yeast. Implications for vacuole biogenesis. *J Biol Chem* 273: 21054–21060.
- Nobrega FG, Nobrega MP & Tzagoloff A (1992) *BCS1*, a novel gene required for the expression of functional Rieske iron-sulfur protein in *Saccharomyces cerevisiae*. *EMBO J* 11: 3821–3829.
- Ohya Y, Umemoto N, Tanida I, Ohta A, Iida H & Anraku Y (1991) Calcium-sensitive *cls* mutants of *Saccharomyces cerevisiae* showing a Pet[−] phenotype are ascribed to defects of vacuolar membrane H⁺-ATPase activity. *J Biol Chem* 266: 13971–13977.
- Patterson TE, Albrecht EB, Nurse P, Sazer S & Stark GR (1999) Effects of genome position and the DNA damage checkpoint on the structure and frequency of *sod2* gene amplification in fission yeast. *Mol Biol Cell* 10: 2199–2208.
- Prior C, Mamessier P, Fukuhara H, Chen XJ & Wésolowski-Louvel M (1993) The hexokinase gene is required for the transcriptional regulation of the glucose transporter gene *RAG1* in *Kluyveromyces lactis*. *Mol Cell Biol* 13: 3882–3889.
- Prior C, Potier S, Souciet JL & Sychrova H (1996) Characterization of the *NHA1* gene encoding a Na⁺/H⁺ antiporter of the yeast *Saccharomyces cerevisiae*. *FEBS Lett* 387: 89–93.


- Sambrook J & Russel DW (2001) *Molecular Cloning: A Laboratory Manual*, 3rd edn. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Serrano R, Marquez JA & Rios G (1997) Crucial factors in salt stress tolerance. *Yeast Stress Responses* (Hohmann S & Mager WH, eds), pp. 147–170. Springer, New York.
- Thompson JD, Higgins DG & Gibson TJ (1994) CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res* **22**: 4673–4680.
- van Bakel H, Strengman E, Wijmenga C & Holstege FC (2005) Gene expression profiling and phenotype analysis of *S. cerevisiae* in response to changing copper reveals six genes with new roles in copper and iron metabolism. *Physiol Genomics* **22**: 356–367.
- Wésolowski-Louvel M & Fukuhara H (1979) The genetic map of transfer RNA genes of yeast mitochondria: correction and extension. *Mol Gen Genet* **170**: 261–275.
- Wésolowski-Louvel M, Tanguy-Rougeau C & Fukuhara H (1988) A nuclear gene required for the expression of the linear DNA-associated killer system in the yeast *Kluyveromyces lactis*. *Yeast* **4**: 71–81.
- Wésolowski-Louvel M, Goffrini P, Ferrero I & Fukuhara H (1992) Glucose transport in the yeast *Kluyveromyces lactis*. I. Properties of an inducible low-affinity glucose transporter gene. *Mol Gen Genet* **233**: 89–96.

Volume 13 Number 8 30 June 1997

yeast

ISSN 0749 503X

YESTE3 13(8) 691-782 (1997)



WILEY

Yeast

Subscriptions

Yeast is published monthly plus one additional issue in March, one in June, one in September and one in December (total 16 issues) by John Wiley & Sons, Ltd.

Subscription Rates

Volume 13 1997 16 issues

Institutional US\$995.00

To subscribe, please contact Journals Subscriptions Department, John Wiley & Sons, Ltd., 1 Oldlands Way, Bognor Regis, West Sussex PO22 9SA, UK
Tel: +44 (0) 1243 779777,
Fax: +44 (0) 1243 843232,
email: cs-journals@wiley.co.uk.

Sample Copies

If you are interested in subscribing, you may obtain a free sample copy by contacting John Wiley & Sons, Ltd. at the above address.

Services

Advertising

To advertise in *Yeast*, contact Caroline Melling, Non-Subscription Sales Manager, John Wiley & Sons, Ltd., Baffins Lane, Chichester, West Sussex PO19 1UD, UK.
Tel: +44 (0) 1243 770351,
Fax: +44 (0) 1243 770429,
email: info-assets@wiley.co.uk

For US-based advertising enquiries contact Susan Levey, Advertising Department, John Wiley & Sons, Inc., 605 Third Avenue, New York, NY 10158-0012, USA,
Tel: +1 (212) 850 8833, Fax: +1 (212) 850 8888.

Mailing Lists

Our high quality, profiled subscription lists may be rented for use with targeted mailing campaigns. For a quotation, contact Caroline Melling at the above UK address.

Reprints

Do you wish to purchase reprints of any article from *Yeast*? Reprints from this journal or any previous issue are available to order. Contact Caroline Melling at the above UK address for details of this service.

Copyright

Copyright © 1997 John Wiley & Sons, Ltd.

All Rights Reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, scanning or otherwise, except as described below, without the permission in writing of the Publisher.

Copying of articles is not permitted except for personal and internal use, to the extent permitted by national copyright law, or under the terms of a licence issued by the national Reproduction Rights Organization (such as Copyright Licensing Agency, 90 Tottenham Court Road, London W1P 9HE, UK or Copyright Clearance Center Inc., 27 Congress Street, Salem, MA 01970, USA). The price for copies made under the CCC or other licences is indicated on the title page of each article. Requests for permission for other kinds of copying, such as copying for general distribution, for advertising or promotional purposes, for creating new collective works or for resale, and other enquiries should be addressed to the Publisher.

Statements and opinions expressed in the articles and communications are those of the individual contributors and not the statements and opinions of John Wiley & Sons, Ltd. Wiley assumes no responsibility or liability for any damage or injury to persons or property arising out of the use of any materials, instructions, methods or ideas contained herein. Wiley expressly disclaims any implied warranties of merchantability or fitness for a particular purpose. If expert assistance is required, the services of a competent professional person should be sought.

Abstracting and Indexing

Yeast is covered by the following abstracting and indexing services: BIOSIS, Biotechnology Citation Index (ISI), CAB Abstracts, Cambridge Scientific Abstracts, Chemical Abstracts Service, Current Awareness in Bio. Sci., Current Contents (CC)—Agriculture, Bio. and Env. Sciences, CC—Life Sciences, Excerpta Medica's Embase, Index Medicus, Publications in Food Microbiology, Reference Update, Research Alert Institute for Scientific Information (ISI), Science Citation Index, SCISEARCH Database (ISI), and Telegen Reporter.

Production Information

Production Details

Printed on acid-free paper.

Typeset, printed and bound by Henry Ling Ltd., Dorchester, Dorset.

The cover photograph shows both *S. cerevisiae* and *S. pombe* yeast cells. *S. cerevisiae* photographs (unpublished) are reproduced with permission of Dr M. Veenhuis, Department of Microbiology, University of Groningen, The Netherlands. *S. pombe* photographs are reproduced with permission of the Imperial Cancer Research Fund, Lincoln's Inn Fields, London.

Identification Statement

Yeast (ISSN 0749-503X/USPS 771-670) is published monthly plus one additional issue in March, one in June, one in September and one in December (total 16 issues) by John Wiley & Sons, Ltd., Baffins Lane, Chichester, West Sussex PO19 1UD, UK. Periodicals postage paid at Rahway, NJ. Air freight and mailing in the USA by Mercury Airfreight International Ltd. Inc., 2323 Randolph Avenue, Avenel, NJ 07001, USA.

USA Postmaster—send address changes to *Yeast*, c/o Mercury Airfreight International Ltd. Inc., 2323 Randolph Avenue, Avenel, New Jersey, NJ 07001, USA.

Secretion of Mouse α -Amylase from *Kluyveromyces lactis*

MASAO TOKUNAGA^{1*}, MATSUJIRO ISHIBASHI¹, DAISUKE TATSUDA¹ AND HIROKO TOKUNAGA¹

¹Laboratory of Applied Microbiology, Faculty of Agriculture, Kagoshima University, 1-21-24, Korimoto, Kagoshima 890, Japan

Received 19 September 1996; accepted 20 November 1996

We constructed two mouse α -amylase secretion vectors for *Kluyveromyces lactis* using the well-characterized signal sequence of the pGKL 128 kDa killer precursor protein. Both *PHO5* and *PGK* expression cassettes from *Saccharomyces cerevisiae* directed the expression of mouse α -amylase in YPD medium at a similar level of efficiency. *K. lactis* transformants secreted glycosylated and non-glycosylated α -amylase into the culture medium and both species were enzymatically active. The *K. lactis*/*S. cerevisiae* shuttle secretion vector pMI6 was constructed, and *K. lactis* MD2/1(pMI6) secreted about four-fold more α -amylase than *S. cerevisiae* YNN27 harboring the same plasmid, indicating that *K. lactis* is an efficient host cell for the secretion and production of recombinant proteins.

© 1997 by John Wiley & Sons, Ltd.

Yeast 13: 699–706, 1997.

No. of Figures: 6. No. of Tables: 2. No. of References: 17.

KEY WORDS — Glycosylation; leader-peptide; signal sequence

INTRODUCTION

Yeast is a very useful host in which to express foreign genes, since it is a safe eukaryotic micro-organism with well-established fermentation technology for large-scale production (Romanos *et al.*, 1992). Since the application of recombinant DNA techniques ranges from medical therapeutics to recombinant enzyme production for food science and for biomass utilization, the low-cost production of useful recombinant proteins has become important in recent biotechnology. A large amount of yeast cells can be easily grown by high density cultivation at lower cost than any other eukaryotic expression system.

We have been investigating highly efficient secretion systems for foreign gene products in *Saccharomyces cerevisiae* (Tokunaga *et al.*, 1987, 1988; Kanaya *et al.*, 1989) and in the fission yeast *Schizosaccharomyces pombe* (Tokunaga *et al.*, 1993). Recently, non-*Saccharomyces* (or alterna-

tive) yeasts such as *Pichia pastoris*, *Hansenula polymorpha* and *Yarrowia lipolytica* have been applied as new host cells (Romanos *et al.*, 1992). Among alternative yeast systems, *Kluyveromyces lactis* has much potential as a host, since it has been used to produce β -galactosidase for the food industry. Furthermore, many types of expression vehicles, such as 2 μ m-like multi-copy pKD1 derivatives, integration and centromeric vectors have been developed (Wésolowski-Louvel *et al.*, 1996). The secretion of high levels of bovine prochymosin (van den Berg *et al.*, 1990), human serum albumin (Fleer *et al.*, 1991a) and human interleukin-1 β (Fleer *et al.*, 1991b) from *K. lactis* cells has been described.

In this study, we describe the efficient expression and secretion of mouse α -amylase from *K. lactis* MD2/1 cells into culture medium. *K. lactis* cells secreted both glycosylated and non-glycosylated α -amylases into the culture medium and both were proven to be enzymatically active by staining after non-denaturing gel electrophoresis. Furthermore,

*Correspondence to: Masao Tokunaga.

Table 1. Description of strains and plasmids used in this study.

Strains and plasmids	Characteristics
<i>Strains</i>	
<i>K. lactis</i>	
MD2/1	<i>MATa, uraA, argA1, lysA1, rag1, rag2, cir⁺</i>
PM6-7A	<i>MATa, uraA, ade2, cir⁺</i>
<i>S. cerevisiae</i>	
YNN27	<i>MATa, trp1, ura3, cir⁺</i>
S150-2B	<i>MATa, his3, leu2, trp1, ura3, cir⁺</i>
W303-1B	<i>MATa, his3, leu2, trp1, ura3, ade2, cir⁺</i>
<i>Plasmids</i>	
pSPHO4	Expression vector for <i>K. lactis</i> using <i>PHO5</i> promoter-terminator cassette of <i>S. cerevisiae</i> and secretion signal from 128 kDa killer precursor protein
pSPGK1	Expression vector for <i>K. lactis</i> using <i>PGK</i> promoter-terminator cassette of <i>S. cerevisiae</i> and secretion signal from 128 kDa killer precursor protein
pSK1	<i>K. lactis</i> / <i>S. cerevisiae</i> shuttle vector
pSMF38TMA	Mouse α -amylase secretion vector for <i>S. cerevisiae</i>
pKA128	Mouse α -amylase secretion vector constructed from pSPHO4
pMI5	Mouse α -amylase secretion vector constructed from pSPGK1
pMI6	Mouse α -amylase secretion vector constructed from pSK1

we directly compared the secretion efficiency of mouse α -amylase from *K. lactis* and *S. cerevisiae* cells using the *K. lactis*/*S. cerevisiae* shuttle vector. We found that *K. lactis* MD2/1 cells harboring the expression shuttle vector pMI6 secreted about four-fold more mouse α -amylase into the culture medium than *S. cerevisiae* YNN27 transformants.

MATERIALS AND METHODS

Yeast strains and medium

The strains and plasmids used in this study are summarized in Table 1. The media were YPD (1% Bacto yeast extract, 2% Bacto peptone and 2% glucose), YP (1% Bacto yeast extract and 2% Bacto peptone) and SC (0.67% yeast nitrogen base, amino acid mixture and 2% glucose; Sherman *et al.*, 1986). YPS (1% Bacto yeast extract, 2% Bacto peptone, 1% starch and 2% agar) plates were used for halo assays of α -amylase activity (Tokunaga *et al.*, 1987). For the tunicamycin treatment of the cells, *K. lactis* transformants were cultured at 30°C in YPD medium containing 0.5 µg/ml of tunicamycin. Plasmids were constructed in *Escherichia coli* HB101. *E. coli* transformants were selected on LB plates containing 100 µg/ml ampicillin.

Construction and transformation of secretion vectors


DNA manipulation was as described by Maniatis *et al.* (1982).

Plasmids were constructed as shown in Figure 1A. The expression and secretion vectors, pSPHO4 and pSPGK1, have been described previously (Wéslowski-Louvel *et al.*, 1996). The plasmid pSMF38TMA (Nishizawa *et al.*, 1987) was digested with *Bam*HI and the 1.4 kb fragment containing the α -amylase gene without a signal sequence was digested with mung-bean nuclease. This fragment was subcloned into the pSPHO4 *Eco*RI site that was digested with mung-bean nuclease and bacterial alkaline phosphatase.

The construction of plasmid pMI5 is also shown in Figure 1A. The same 1.4 kb fragment containing the α -amylase gene was subcloned into the *Eco*RI site of pSPGK1, that was digested with mung-bean nuclease and bacterial alkaline phosphatase.

Plasmid pMI5 was digested with *Bam*HI and *Sal*II to isolate a fragment containing the *PGK* promoter, the signal sequence of the 128 kDa killer precursor protein, the mouse α -amylase gene without its own signal sequence and the *PGK*

A.



B.




Figure 1. Construction of secretion plasmids pKA128 and pMI5. (A) Details are described in Materials and Methods. Expression vectors, pSPHO4 and pSPGK1, have been described previously (Wésolowski-Louvel *et al.*, 1996). PHO5-P, promoter of *PHO5*; S.S., signal sequence of the 128 kDa killer precursor protein; PHO5-T, terminator of *PHO5*; S11, replication origin of pKD1; AMP, ampicillin resistance; Amy, mouse α -amylase; PGK-P, promoter of *PGK*; PGK-T, terminator of *PGK*; BAP, bacterial alkaline phosphatase. (B) N-terminal sequence of α -amylase in secretion vectors, pKA128, pMI5 and pMI6. Double underline shows signal sequence derived from the gene encoding 128 kDa killer precursor protein. Underline shows the linker sequence and the thick arrow represents the α -amylase gene. The 19th Pro residue in the present secretion vector corresponds to the 19th residue of mouse salivary α -amylase precursor protein.

transcription termination signal. This fragment was subcloned into *Bam*HI-*Sal*I-digested pSK1 (the *K. lactis*/*S. cerevisiae* shuttle vector; Prior *et al.*, 1993) to construct pMI6.

The yeast was transformed using the Li^+ -salt described by Ito *et al.* (1983).

Assay of α -amylase activity

For the halo assay, yeast transformants were patched onto YPS plates and incubated for 48–72 h at 30°C. Halos surrounding the transformants were detected by the absence of iodo-staining (I_2 -KI) of the digested starch (Tokunaga *et al.*, 1993).

The activity of mouse α -amylase secreted into the medium was determined as described by Nelson (1944) and Somogyi (1952). Transformants were precultured in SC medium and 0.5 ml of this seed culture was inoculated onto 5 ml of rich medium (YP plus carbon source) for the production and secretion of α -amylase. The reaction mixture containing 0.5 ml of 1% soluble starch dissolved in 15 mM-N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid (HEPES) buffer (pH 7.0) with 1.5 mM- CaCl_2 and 0.5 ml of crude enzyme (culture supernatant dialysed against 15 mM-HEPES buffer, pH 7.0) were incubated at 37°C. At the appropriate time, 0.2 ml of reaction mixture was mixed with the Somogyi solution, then the Nelson solution, as described (Nelson, 1944; Somogyi, 1952).

Active staining of α -amylase on non-denatured polyacrylamide gel electrophoresis (PAGE) proceeded as follows. Crude enzyme was resolved on 6.5% polyacrylamide non-denatured gels with 25 mM-Tris/192 mM-glycine buffer (pH 8.3) at 20 mA for 6 h at 4°C. The gel after electrophoresis was incubated in 50 mM-HEPES buffer (pH 7.0) containing 0.02% soluble starch for 1 h at room temperature, then stained with 0.2% KI-2% I_2 .

Sodium dodecyl sulfate-polyacrylamide gel electrophoresis and immunoblotting

Proteins were separated by sodium dodecyl sulfate (SDS)-PAGE as described by Laemmli (1970). Proteins resolved on SDS-PAGE were electroblotted onto a nitrocellulose membrane using a semi-dry blotting apparatus (BIO CRAFT BE-300) and a buffer of 25 mM-Tris/192 mM-glycine/20% methanol, pH 8.3. The blotted α -amylase was immunostained with an anti-human- α -amylase IgG/peroxidase-conjugated anti-rabbit-IgG antibody/4-chloronaphthol- H_2O_2 (Tokunaga *et al.*, 1983, 1987, 1988).

RESULTS AND DISCUSSION

Expression and secretion of mouse α -amylase from *K. lactis* transformants

We constructed α -amylase secretion vectors using the signal peptide of pGKL killer 128 kDa precursor protein and multi-copy expression vectors (Figure 1). This signal sequence has been applied to *S. cerevisiae* (Tokunaga *et al.*, 1987), *Sz. pombe* (Tokunaga *et al.*, 1993) and *K. lactis* (Fleer *et al.*, 1991a,b) recombinant systems.




Figure 2. Halo assay and immunoblotting of secreted mouse α -amylase. (a) Halo assay of pKA128 and pMI5. Controls without the α -amylase gene are also shown. (b) Immunoblotting of secreted mouse α -amylase. Proteins in the culture supernatant (1.5 ml) were concentrated by 10% trichloroacetic acid precipitation, resolved by SDS-PAGE, then immunoblotted. Lane 1, pre-stained marker; lane 2, *K. lactis* (pSPHO4); lane 3, *K. lactis* (pKA128); lane 4, *K. lactis* (pSPGK1); lane 5, *K. lactis* (pMI5); lane 6, authentic human salivary α -amylase (Sigma IX-A), the antigen used for preparing anti- α -amylase antibody.

The secretion of mouse α -amylase from *K. lactis* MD2/1 cells is shown in Figure 2. *K. lactis* cells harboring plasmids pKA128 and pMI5 formed clear halos on YPS plates, indicating that *K. lactis* can secrete α -amylase into the culture medium using the signal peptide of the pGKL killer 128 kDa precursor protein (Figure 2a). The size of the *K. lactis* (pKA128) halo was similar to that of *K. lactis* (pMI5), suggesting that *PHO5* and *PGK* expression cassettes of *S. cerevisiae* function sufficiently well to support good levels of secretion when used with a suitable leader sequence in *K. lactis* transformants. *K. lactis* harboring the vector plasmid pSPHO4 and pSPGK1 without the α -amylase gene did not produce halos. We also constructed α -amylase secretion vectors using the signal peptide of pGKL killer 28 kDa precursor protein, but the efficiency of secretion was lower




Figure 3. Immunoblotting and active staining of secreted mouse α -amylases with and without tunicamycin. (a) Immunoblotting of α -amylase. Lane 1, prestained marker; culture with (lane 3) and without (lane 2) 0.5 μ g/ml tunicamycin. (b) Active staining of α -amylases. *K. lactis* (pMI5) cultured with (lane 1) and without (lane 2) tunicamycin. Lane 3, authentic human salivary α -amylase.

than that of the 128 kDa protein (data not shown) as found in *Sz. pombe* transformants (Tokunaga *et al.*, 1993).

We identified α -amylase secreted into the culture medium by means of Western blotting using anti-human- α -amylase antibody. As shown in Figure 2b, culture supernatants of *K. lactis* (pKA128) and *K. lactis* (pMI5) contained two bands with a molecular mass of around 55 000, which cross-reacted with anti-human α -amylase antibody (lanes 3 and 5). No bands were evident in *K. lactis* cells containing pSPHO4 or pSPGK1 (lanes 2 and 4).

Glycosylation of secreted α -amylase

It has been reported that recombinant mouse α -amylase secreted into the culture medium by *S. cerevisiae* is composed of glycosylated and non-glycosylated molecules (Tokunaga *et al.*, 1988, 1992). To clarify the glycosylation status of α -amylase, *K. lactis* cells were grown in the presence of tunicamycin (0.5 μ g/ml culture) at 30°C overnight. The α -amylase secreted into the culture medium was analysed by SDS-PAGE and immunoblotting. As shown in Figure 3a, the upper band disappeared in the presence of tunicamycin (lane 3). These findings indicated that the upper and lower bands are glycosylated and non-glycosylated α -amylase, respectively, as seen in the *S. cerevisiae* secretion system (Tokunaga *et al.*, 1987, 1992).

To determine whether or not both glycosylated and non-glycosylated α -amylase molecules are en-




Figure 4. Effects of carbon sources on α -amylase secretion. Immunoblotting of secreted mouse α -amylase. Proteins in the culture supernatant (1.5 ml) were concentrated by 10% trichloroacetic acid precipitation, then resolved by SDS-PAGE and immunoblotted. The final cell growth (Klett value) is also shown.

zymatically active, we actively stained α -amylase activity on non-denaturing PAGE. Figure 3b, lanes 1 and 2, shows that both glycosylated and non-glycosylated α -amylases were enzymatically active. The upper band was glycosylated α -amylase and the lower was non-glycosylated α -amylase on non-denatured PAGE, since the former disappeared in the presence of tunicamycin (lane 1). We are the first to demonstrate that both glycosylated and non-glycosylated recombinant mouse salivary α -amylases are enzymatically active. In a control experiment, authentic human α -amylase (Sigma IX-A) migrated much faster than recombinant mouse α -amylases in this non-denaturing gel system. The upper glycosylated band that was bound to concanavalin A-Sepharose (data not shown) and the lower non-glycosylated band were both enzymatically active (lane 3).

Effects of carbon sources on the efficiency of α -amylase production

One advantage of *K. lactis* compared with *S. cerevisiae* is that it can utilize a wide variety of carbon sources (Wésolowski-Louvel *et al.*, 1996). For example, *K. lactis* assimilates lactose, L-sorbose, and L-lysine as carbon sources, but *S. cerevisiae* does not. We examined the effect of carbon sources on α -amylase secretion. Various carbon sources were added to YP medium, then we examined the cell growth of MD2/1 (pMI5) and secretion of α -amylase (Figure 4).

When glucose, lactose or galactose was used as a carbon source, the enzyme was efficiently secreted. When glycerol, ethanol or lysine was the carbon




Figure 5. Comparison of the efficiency of α -amylase secretion between *K. lactis* and *S. cerevisiae* using the *K. lactis*/*S. cerevisiae* shuttle secretion vector, pMI6. (a) Halo assay of α -amylase secretion; *S. cerevisiae* YNN27 (pMI6) and *K. lactis* MD2/1(pMI6) are shown. (b) Immunoblots of the same samples. Trichloroacetic acid precipitates of culture medium (1.5 ml) were analysed.

source, the secretion gradually decreased, and the levels were very low in the presence of L-sorbose.

*Direct comparison of secretion efficiency of α -amylase from *K. lactis* and *S. cerevisiae* using the *K. lactis*/*S. cerevisiae* shuttle secretion vector pMI6*

The *K. lactis*/*S. cerevisiae* shuttle vector, pSK1, has replication origins from *K. lactis* (S11 fragment of pKD1) and *S. cerevisiae* (*ori* of 2 μ m) plasmids. This shuttle plasmid was originally constructed by Prior *et al.* (1993) to characterize *K. lactis* homologous gene complementing an *S. cerevisiae* mutation and *vice versa*. Here, we constructed the shuttle secretion vector, pMI6, by inserting a mouse α -amylase secretion cassette encoded by the 3.2 kb *Sall*-*Bam*HI fragment of pMI5 (*PGK* promoter-signal sequence of the 128 kDa killer precursor- α -amylase-*PGK* terminator) into *Sall*/*Bam*HI-digested pSK1. Two strains of *K. lactis*, MD2/1 and PM6-7A, and three strains of *S. cerevisiae*, YNN27, S150-2b and W303-1B, were transformed with pMI6 in such a manner that the efficiency of α -amylase secretion from *K. lactis* and *S. cerevisiae* cells can be directly compared using the same secretion vector. These strains have been used often to study the secretion of heterologous proteins (Tokunaga *et al.*, 1990; Wésolowski-Louvel *et al.*, 1996).

Table 2. α -Amylase activities secreted from *K. lactis* and *S. cerevisiae* cells harboring *K. lactis*/*S. cerevisiae* shuttle secretion vector pMI6.

Strains	α -Amylase activity (μ mol/min per ml medium)	Final cell density (Klett value)
<i>K. lactis</i>		
MD2/1	0.527	560
PM6-7A	0.148	630
<i>S. cerevisiae</i>		
YNN27	0.141	550
S150-2B	0.006	455
W303-1B	0.009	540

Transformants were initially cultured in SC medium and 0.5 ml of this seed culture was inoculated onto 5 ml of YP-2% galactose medium. After 72 h, activity secreted into the medium was assayed.

As shown in Figure 5a, both transformants formed clear halos, indicating that this shuttle vector functioned in both strains. *K. lactis* MD2/1 (pMI6) formed a larger halo than that of *S. cerevisiae* YNN27(pMI6). The immunoblotting profile was also the same, as shown in Figure 5b. The α -amylase activities in the culture supernatants assayed as described by Somogyi and Nelson revealed that *K. lactis* MD2/1(pMI6) secretes about four-fold more α -amylase activity than *S. cerevisiae* YNN27(pMI6) (Table 2). *K. lactis* PM6-7A(pMI6), which secretes less than MD2/1, secreted almost the same amount of α -amylase as *S. cerevisiae* YNN27(pMI6). Other *S. cerevisiae* strains harboring pMI6 secreted much less α -amylase; we could not resolve the significant difference in the secretion amounts among *S. cerevisiae* strains. These data indicated that *K. lactis* is more suitable as a host than *S. cerevisiae* with respect to the secretion of recombinant mouse α -amylase.

We measured the plasmid stability of MD2/1 (pMI6) and YNN27(pMI6) under the same conditions as the α -amylase assay described above. After 72 h of culture in rich medium without selective pressure, the stability of pMI6 was 11.4% in *K. lactis* MD2/1 cells, and 22.0% in *S. cerevisiae* YNN27 cells (average value in four experiments). *K. lactis* still secreted much more α -amylase regardless of its lower capacity for plasmid distribution than *S. cerevisiae*. This indicated that the increased plasmid stability in *K. lactis* cells will much improve the secretion efficiency.




Figure 6. Cell growth of yeast transformants in the presence of starch. *K.l.*, *K. lactis* MD2/1(pMI6); *S.c.*, *S. cerevisiae* YNN27(pMI6); (+), YP medium containing 2% soluble starch; (-), YP medium without starch.

Assimilation of starch by *K. lactis* transformants

To determine whether or not *K. lactis* transformant cells secreting α -amylase can utilize starch as a carbon source for growth, *K. lactis* MD2/1(pMI6) and *S. cerevisiae* YNN27(pMI6) were cultured in YP medium with and without 2% soluble starch. As shown in Figure 6, the amount of *K. lactis* growth in YP medium with 2% starch was about two-fold higher than that without starch. The growth level of *K. lactis* transformants with starch was similar to that in YPD medium. The growth of *S. cerevisiae*, which was one-third that of *K. lactis* without starch, did not differ in YP medium with or without 2% starch. These data indicated that although both *K. lactis* and *S. cerevisiae* cells partly utilized the ingredients of YP medium as carbon sources, the former can efficiently assimilate oligosaccharides derived from starch digested by secreted α -amylase and can thus support their own growth. In contrast, *S. cerevisiae* cells harboring mouse α -amylase could not utilize oligosaccharides derived from starch as the carbon source (Figure 6). It is also conceivable that the total amount of α -amylase production from *S. cerevisiae* cells was lower than that from *K. lactis* cells under growth conditions using YP or YP-starch as the carbon source.

In conclusion, we induced the secretion of mouse α -amylase into the culture medium of *K. lactis* transformants using the pGKL 128 kDa

killer secretion signal, and showed that *K. lactis* is superior to *S. cerevisiae* and *Sz. pombe* (Tokunaga *et al.*, 1993) for producing recombinant mouse α -amylase.

ACKNOWLEDGEMENT

We are grateful to Dr Hiroshi Fukuhara of Institut Curie for the gift of strains and plasmids, and for helpful comments regarding the manuscript.

REFERENCES

- van den Berg, J. A., van der Laken, K. J., van Ooyen, A. J. J., *et al.* (1990). *Kluyveromyces* as a host for heterologous gene expression: expression and secretion of prochymosin. *Biol/Technology* **8**, 135–139.
- Fleer, R., Yeh, P., Amellal, N., *et al.* (1991a). Stable multicopy vectors for high-level secretion of recombinant human serum albumin by *Kluyveromyces lactis*. *Biol/Technology* **9**, 968–975.
- Fleer, R., Chen, X. J., Amellal, N., *et al.* (1991b). High-level secretion of correctly processed recombinant human interleukin-1 β in *Kluyveromyces lactis*. *Gene* **107**, 285–295.
- Ito, H., Fukuda, Y., Murata, K. and Kimura, A. (1983). Transformation of intact yeast cells treated with alkali cations. *J. Bacteriol.* **153**, 163–168.
- Kanaya, E., Higashizaki, T., Ozawa, F., *et al.* (1989). Synthesis and secretion of human nerve growth factor by *Saccharomyces cerevisiae*. *Gene* **83**, 65–74.
- Laemmli, U. K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**, 680–685.
- Maniatis, T., Fritsch, E. F. and Sambrook, J. (1982). *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory, Cold Spring Harbor, New York.
- Nelson, N. (1944). A photometric adaptation of the Somogyi method for the determination of glucose. *J. Biol. Chem.* **153**, 375–380.
- Prior, C., Mamessier, P., Fukuhara, H., Chen, X. J. and Wesołowski-Louvel, M. (1993). The hexokinase gene is required for transcriptional regulation of the glucose transporter gene *RAG1* in *Kluyveromyces lactis*. *Mol. Cell. Biol.* **13**, 3882–3889.
- Romanos, M. R., Scorer, C. A. and Clare, J. J. (1992). Foreign gene expression in yeast: a review. *Yeast* **8**, 423–488.
- Sherman, F., Fink, G. R. and Hicks, J. B. (1986). *Laboratory Course Manual for Methods in Yeast Genetics*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, pp. 163–167.
- Somogyi, M. (1952). Notes on sugar determination. *J. Biol. Chem.* **195**, 19–23.
- Tokunaga, M., Wada, N. and Hishinuma, F. (1987). A novel yeast secretion vector utilizing secretion signal of killer toxin encoded on the yeast linear DNA

- plasmid pGKL1. *Biochem. Biophys. Res. Commun.* **144**, 613–619.
- Tokunaga, M., Wada, N. and Hishinuma, F. (1988). A novel secretion signal isolated from 28K killer precursor protein encoded on the linear DNA plasmid pGKL1. *Nucl. Acids Res.* **15**, 7499–7511.
- Tokunaga, M., Kawamura, A., Omori, A. and Hishinuma, F. (1992). Characterization of mouse α -amylase secreted from *Saccharomyces cerevisiae* by the pGKL 128 kDa killer secretion signal. *Biosci. Biotech. Biochem.* **56**, 155–156.
- Tokunaga, M., Kawamura, A., Yonekyuu, S., Kishida, M. and Hishinuma, F. (1993). Secretion of mouse α -amylase from fission yeast *Schizosaccharomyces pombe*: presence of chymostatine-sensitive protease activity in the culture medium. *Yeast* **9**, 379–387.
- Wésolowski-Louvel, M., Breunig, K. and Fukuhara, H. (1996). *Kluyveromyces lactis*. In Wolf, K. (Ed.), *Non-Conventional Yeasts in Biotechnology, a Handbook*. Springer-Verlag, Berlin, Heidelberg, p. 139–201.

Current Genetics

33/1

98

Eukaryotes
with Emphasis on
Yeasts
Fungi
Protists
Cell Organelles



Springer

Current Genetics

Eukaryotes
Yeasts · Fungi · Protists · Cell Organelles

Synopsis

Current Genetics is an international journal devoted to the rapid publication of original articles of immediate importance on genetics of eukaryotes, with emphasis on yeasts, other fungi, protists and cell organelles. Papers on formal genetics will only be accepted if they contribute to the understanding of special genetic systems or give access to molecular approaches

Copyright

Submission of a manuscript implies: that the work described has not been published before (except in the form of an abstract or as part of a published lecture, review, or thesis); that it is not under consideration for publication elsewhere; that its publication has been approved by all coauthors, if any, as well as by the responsible authorities at the institute where the work has been carried out; that, if and when the manuscript is accepted for publication, the authors agree to automatic transfer of the copyright to the publisher; and that the manuscript will not be published elsewhere in any language without the consent of the copyright holders.

All articles published in this journal are protected by copyright, which covers the exclusive rights to reproduce and distribute the article (e.g., as offprints), as well as all translation rights. No material published in this journal may be reproduced photographically or stored on microfilm, in electronic data bases, video disks, etc., without first obtaining written permission from the publisher.

The use of general descriptive names, trade names, trademarks, etc., in this publication, even if not specifically identified, does not imply that these names are not protected by the relevant laws and regulations.

While the advice and information in this journal is believed to be true and accurate at the date of its going to press, neither the authors, the editors, nor the publisher can accept any legal responsibility for any errors or omissions that may be made. The publisher makes no warranty, express or implied, with respect to the material contained herein.

Special regulations for photocopies in the USA: Photocopies may be made for personal or in-house use beyond the limitations stipulated under Section 107 or 108 of U.S. Copyright Law, provided a fee is paid. All fees should be paid to the Copyright Clearance Center, Inc., 21 Congress Street, Salem, MA 01970, USA, stating the ISSN 0172-8083, the volume, and the first and last page numbers of each article copied. The copyright owner's consent does not include copying for general distribution, promotion, new works, or resale. In these cases, specific written permission must first be obtained from the publisher.

The Canada Institute for Scientific and Technical Information (CISTI) provides a comprehensive, world-wide document delivery service for all Springer-Verlag journals. For more information, or to place an order for copyright-cleared Springer-Verlag document, please contact Client Assistant, Document Delivery, Canada Institute for Scientific and Technical Information, Ottawa K1A 0S2, Canada (Tel: 613-993-9251; Fax: 613-952-8243; e-mail: cisti-dodcel@nrc.ca).

Subscription information

ISSN print edition 0172-8083
ISSN electronic edition 1432-0983
Volumes 33 and 34 (6 issues each) will appear in 1998.

North America. Recommended annual subscription rate: approx. US \$ 1448.00 (single issue price: approx. US \$ 140.60) including carriage charges. Subscriptions are entered with prepayment only. Orders should be addressed to:
Springer-Verlag New York Inc., Journal Fulfillment Services Department, P.O. Box 2485 Secaucus, NJ 07096-2485, USA
Tel. +1-800-Springer; Fax +1-201-348 4505

All other countries. Recommended annual subscription rate: DM 2236.00 plus carriage charges; [Germany DM 34.80, incl. value added tax; all other countries: DM 50.40 except for the following countries to which SAL delivery (Surface Airmail Lifted) is mandatory: Japan, India, Australia/New Zealand. SAL charges and airmail delivery to all other countries are available upon request].
Volume price: DM 1118.00, single issue price: DM 223.60 plus carriage charges. Orders can either be placed via a bookdealer or sent directly to:
Springer-Verlag, Karin Tiks, Heidelberger Platz 3 D-14197 Berlin, Germany
Tel. +49-30-82787-358; Fax +49-30-82787-448

Cancellations must be received by September 30 to take effect at the end of the same year.

Changes of address. Allow six weeks for all changes to become effective. All communications should include both old and new addresses (with postal codes) and should be accompanied by a mailing label from a recent issue.

According to § 4 section 3 of the German Postal Services Data Protection Regulations, the German Federal Post Office can inform the publisher of a subscriber's new address even if the subscriber has not submitted a formal application for mail to be forwarded.

Subscribers not in agreement with this procedure may send a written complaint to Springer-Verlag's Berlin office within 14 days of publication of this issue.

Back volumes. Prices are available on request.

Microform editions are available from:
University Microfilms International
300 N. Zeeb Road, Ann Arbor, MI 48106, USA

Electronic edition. An electronic edition of this journal is available via the LINK Information Service <http://link.springer.de>. You may contact us per e-mail access@link.springer.de or by Fax +49-6221-487-288. Would you like to automatically receive per e-mail the table of contents with direct links to the respective abstracts as soon as a new issue is available via LINK? Then take advantage of our new, free service LINK Alert. For registration and further information just go to <http://link.springer.de/alert>

Production

Springer-Verlag, May Doebl
Journal Production Life Sciences
Postfach 10 52 80, D-69042 Heidelberg, Germany
Tel. +49-6221-487959; Fax +49-6221-487967
e-mail: doebl@springer.de

Address for courier, express and registered mail: Tiergartenstrasse 17, D-69121 Heidelberg, Germany

Responsible for advertisements

Springer-Verlag, E. Lückermann
Heidelberger Platz 3, D-14197 Berlin, Germany
Tel. +49-30-82787-0; Fax +49-30-82787-300

Printers

Konrad Tritsch, GmbH, Würzburg, Germany
© Springer-Verlag Berlin Heidelberg 1998
Springer-Verlag GmbH & Co. KG
Berlin, Germany
Printed in Germany



Springer

ORIGINAL PAPER

Philip M. Hansbro · Xin Jie Chen
G. Desmond Clark-Walker

Allele-specific expression of the Mgi^- phenotype on disruption of the F_1 -ATPase delta-subunit gene in *Kluyveromyces lactis*

Received: 9 June / 27 September 1997

Abstract *Kluyveromyces lactis* is a petite-négative yeast that does not form viable mitochondrial genome-deletion mutants (petites) when treated with DNA-targeting drugs. Loss of mtDNA is lethal for this yeast but mutations at three loci termed *MGI*, for mitochondrial genome integrity, can suppress this lethality. The three loci encode the α -, β - and γ -subunits of mitochondrial F_1 -ATPase. In this study we report the isolation and characterization of the *KIATP δ* gene encoding the δ -subunit of F_1 -ATPase. The deduced protein contains 158 amino acids showing 72% identity to the protein from *Saccharomyces cerevisiae* and a putative mitochondrial targeting sequence of 23 amino acids. Disruption of the gene causes cells to become respiratory deficient while the introduction of *ATP δ* from *S. cerevisiae* restores growth on glycerol. Cells with a disrupted *ATP δ* gene, like strains with disruptions of α -, β - and γ - F_1 -subunits, do not produce petite mutants when treated with ethidium bromide. However, unlike strains with disruptions in the three largest F_1 -subunits, disruption of *ATP δ* in the presence of some *mgi* alleles does not abolish the Mgi^- phenotype. By contrast, elimination of *ATP δ* in other *mgi* strains removes resistance to ethidium bromide and ρ^0 mutants are not formed. Hence the *ATP δ* subunit of F_1 -ATPase, while not mandatory for a Mgi^- phenotype, aids some *mgi* alleles in suppressing ρ^0 lethality.

Key words *Kluyveromyces lactis* · Mitochondrial genome integrity · F_1 -ATPase δ -subunit gene · *ATP δ* -disruption

P. M. Hansbro¹ · X. J. Chen · G. D. Clark-Walker (✉)
Molecular and Cellular Genetics Group,
Research School of Biological Sciences,
The Australian National University, PO Box 475,
Canberra City, ACT, 2601, Australia,
Fax No: +61-6-279 8294,
e-mail: dcw@rsbs-central.anu.edu.au

Present address:

¹ Department of Pathology, University of Cambridge,
Cambridge, CB2 6PD, UK

Communicated by R. J. Schweyen

Introduction

The energy transducing $F_1 F_0$ -ATP synthases of bacteria, chloroplasts and mitochondria use a proton gradient, generated across a membrane by respiration or photosynthesis, to synthesize ATP (Futai et al. 1989; Senior 1990; Cox et al. 1992; Hatefi 1993; Boyer 1993; Deckers-Hebestreit and Altendorf 1996). In mitochondria, the F_0 component, containing 8–11 different subunits, spans the inner membrane, while the F_1 complex, composed of five proteins with the stoichiometry 3α , 3β , 1γ , 1δ , 1ϵ , extends into the matrix (Attardi and Schatz 1988; Grivell 1989; Poyton and McEwen 1996). For mitochondrial ATP-synthase, genes for the five F_1 subunits are encoded by nuclear DNA whereas genes for either two or three of the 8–11 subunits of the F_0 complex are located in mitochondrial DNA (mtDNA) (Poyton and McEwen 1996). Although the F_1 component of bacteria, chloroplasts and mitochondria all contain five separate subunits, only those of bacteria and chloroplasts share the same nomenclature and sequence homology (Cox et al. 1992). Differences between mitochondria and bacteria are found in the two smallest F_1 subunits. In bacteria the ϵ -subunit shares sequence homology with the δ -subunit of mitochondria while the ϵ -subunit of mitochondria does not have an equivalent protein in bacteria.

A better understanding of the role played by the three F_1 subunits in the operation of ATP synthase has been provided by structural analysis of bovine mitochondrial F_1 (Abrahams et al. 1994). The α - and β -subunits alternate in a hexameric complex where the catalytic site for ATP synthesis occurs in β -subunits at a contact region with α -subunits. A central space formed by the hexameric array is occupied by amino- and carboxy terminal α -helices of the γ -subunit. Apart from another small portion of the γ -subunit, the central part of this protein, together with the δ - and ϵ -subunits, have not been located. One view is that these proteins form part of a stalk that helps F_1 attach to F_0 in the membrane (Walker and Collinson 1994; Deckers-Hebestreit and Altendorf 1996).

Table 1 Genotype and source of yeast strains

Strain	Relevant genotype	Source or reference
CK56-7A	α , <i>ade1</i> , <i>lysA1</i> , <i>uraA1</i> , <i>mgil-1</i>	Chen and Clark-Walker (1996)
CK99-7C	α , <i>his7</i> , <i>metA1</i> , <i>trpX</i> , <i>uraA1</i> , <i>mgil2-1</i> , <i>mex1-1</i>	Chen and Clark-Walker (1995)
CK103-1B	α , <i>ade1</i> , <i>adeT-600</i> , <i>lysA1</i> , <i>uraA1</i> , <i>mgil-9</i>	Clark-Walker et al., in preparation
CK113/1	α , <i>ade1</i> , <i>uraA1</i> , <i>mgil2-6</i>	Clark-Walker et al., in preparation
CK359/1	α , <i>ade1</i> , <i>lysA1</i> , <i>uraA1</i> , <i>mgil-1</i> , <i>atpδ::URA3</i>	This study
CK361/2	α , <i>his7</i> , <i>metA1</i> , <i>trpX</i> , <i>uraA1</i> , <i>mgil2-1</i> , <i>mex1-1</i> , <i>atpδ::URA3</i>	This study
CK362/1	α , <i>ade1</i> , <i>uraA1</i> , <i>mgil2-6</i> , <i>atpδ::URA3</i>	This study
CK363/1	α , <i>metA1</i> , <i>uraA1</i> , <i>mgil5-2</i> , <i>atpδ::URA3</i>	This study
CK364/1	α , <i>ade1</i> , <i>ade2</i> , <i>lysA1</i> , <i>uraA1</i> , <i>mgil-9 atpδ::URA3</i>	This study
CW26-1A	α , <i>metA1</i> , <i>uraA1</i> , <i>mgil5-2</i>	Clark-Walker et al., in preparation
PM6-7A	α , <i>ade2</i> , <i>uraA1</i>	Chen <i>et al.</i> (1992)
PM67A/ <i>atpδ::URA3</i>	α , <i>ade2</i> , <i>uraA1</i> , <i>atpδ::URA3</i>	This study

Evidence from parallel studies with *Escherichia coli* and spinach chloroplasts supports the view that the γ -subunit rotates relative to the hexameric ring of the α - and β -subunits (Duncan et al. 1995; Sabbert et al. 1996; Noji et al. 1997) and that in *E. coli* the ε -subunit (equivalent to the δ -subunit in mitochondria) shows relative movement during ATP hydrolysis (Aggeler and Capaldi 1996).

Our interest in the yeast mitochondrial δ -subunit has arisen from investigations into a phenomenon seemingly unrelated to ATP synthesis. A minority of yeasts, exemplified by *Saccharomyces cerevisiae*, readily form respiratory deficient (petite) mutants that lose mtDNA when treated with DNA targeting drugs (Ephrussi 1953; Bulder 1964 a, b; De Deken 1966; Slonimski et al. 1968; Clark-Walker et al. 1981). By contrast, *Kluyveromyces lactis*, as a representative of most yeasts, does not form viable petite-negative mutants upon exposure to such drugs (Bulder 1964 a b; Heritage and Whittaker 1977). Indeed, elimination of mtDNA is lethal for wild-type *K. lactis* (Clark-Walker and Chen 1996). However, it has been discovered using this yeast that a class of nuclear genome mutants termed *mgil*, for mitochondrial genome integrity, can suppress the petite-negative phenotype so that mtDNA deletion mutants can be recovered (Chen and Clark-Walker 1993). Characterization of three *MGI* loci has shown that they encode the α -, β - and γ -subunits of F_1 -ATPase (Chen and Clark-Walker 1995, 1996). Specific mutations, producing amino-acid substitutions, together with an assembled F_1 particle, appear to be necessary for a *Mgi*⁻ phenotype as disruption of any of these three genes results in the loss of suppressor activity.

As a foundation for the possible identification of further *MGI* loci, and to gain an understanding of how *mgil* mutations permit the loss of mtDNA, we undertook to isolate and characterize the *ATP δ* gene encoding the F_1 δ -subunit of *K. lactis*, and to examine any possible involvement of this subunit in the conversion between petite-negative and -positive phenotypes.

Materials and methods

Strains and media. The *K. lactis* strains used in this study are listed in Table 1. The isolation and characterization of three new *mgil* alleles *mgil-9*, 2-6 and 5-2, not previously described, will be reported in a subsequent publication (Clark-Walker et al., in preparation). *S. cerevisiae* AH22 *Mata*, *leu2-3, 112*, *his4* was used as a source of genomic DNA for PCR amplification. Complete medium (GYP) contains 0.5% Bacto yeast extract, 1% Bacto Peptone, and 2% glucose. Glycerol medium (GlyYP) contains 2% glycerol in place of glucose. Minimal medium (GMM) contains 0.67% Difco yeast nitrogen base without amino acids and 2% glucose. Nutrients essential for auxotrophic strains were added at 25 μ g/ml for bases and 50 μ g/ml for amino acids. Ethidium bromide (EB) medium is GYP plus EB at 16 μ g/ml.

Amplification by the polymerase chain reaction. Amplification of *ATP δ* from *S. cerevisiae* was carried out as described by Mullis and Faloona (1987) using a *Taq* DNA polymerase kit supplied by Promega. The forward primer, 5'(+134)GTGGCTCCGAAGTTACTC(+151)3', and reverse primer, 5'(449)TCTACTTGAATGACAGC(433)3', were designed from the *S. cerevisiae ATP δ* sequence (Giraud and Velours 1994) and incorporated into a reaction with 50 ng of DNA from AH22 (Chen and Clark-Walker 1995). The purified PCR product had an electrophoretic mobility equivalent to 316-bp of the expected fragment.

Isolation and sequencing of *K. lactis ATP δ* . The PCR-amplified fragment of *ATP δ* was labelled with [³²P] dATP by the random primer method and used to probe a *K. lactis* partial *Sau3A1* gene bank cloned in the *K. lactis/E. coli* shuttle vector KEp6 (Weslowski-Louvel et al. 1986). The two positive colonies were subjected to further analysis and one, containing a plasmid with a 5-kb insert, was mapped using restriction endonucleases. For nucleotide-sequence determination, appropriate DNA fragments were cloned into the sequencing vectors pTZ18 U and pTZ19 U (Pharmacia) and single-stranded templates were subsequently obtained for sequence determination using the dideoxy chain-termination procedure (Sanger et al. 1977). In addition to the 'universal primer' a synthetic oligonucleotide, 5'(+287)CTGTCCAACCAGACT(+301)3', was prepared to complete the sequence.

Disruption of *ATP δ* . Gene disruption was carried out by the one-step replacement procedure (Rothstein 1983). A plasmid containing a 1.65-kb *StuI-BamHI* fragment (see Fig. 1), was opened at the unique *BglII* site within the *ATP δ* gene and a 1.1-kb *BglII* fragment containing the *URA3* gene of *S. cerevisiae* was inserted. Disruption of *ATP δ*

in *K. lactis* PM6-7A and various *mg1* strains was achieved by transformation with a 1.74-kb *MscI*-*BspHI* fragment obtained from the above construct. Transformants were selected for *Ura*⁺ and the correct gene replacement was verified by digestion of genomic DNA with *MscI*-*BspHI* and hybridization to [³²P]-labelled *ATPδ*.

Complementation of *K. lactis* containing a disrupted *ATPδ*. The plasmid pCXJ22-*ScATPδ* was constructed by cloning a 2.6-kb *Bam*HI-*Eco*RI fragment, containing *ScATPδ* isolated from pMFG4 (Giraud and Velours 1994), into the *Bam*HI and *Eco*RI sites of the *K. lactis* vector pCXJ22 (Chen 1996). The *KIATPδ*-disrupted strain PM6-7A/*ATPδ*::*URA3* was transformed with pCXJ22-*ScATPδ*. Transformed cells were plated on GMM plates, grown at 28°C for 24 h, and then replica-plated to GlyYP medium. Respiratory competent colonies (Gly⁺) were obtained, indicating that *ScATPδ* can complement the disrupted allele of *KIATPδ*.

Viability determination. Cells were grown in GYP medium to stationary phase and spread onto the surface of a GYP agar slice. Individual cells were removed with a de Fonbrun micromanipulator to fresh medium to allow the development of individual colonies.

Nucleotide-sequence accession number. GenBank U88046. The sequence of the *K. lactis* F₁-ATPase δ -subunit gene, *KIATPδ* has been given the GenBank accession number U88046.

Results and discussion

Cloning and sequencing of *ATPδ*

A *K. lactis* gene bank was probed with a fragment of the δ -subunit gene from *S. cerevisiae*. Of the two positive clones identified, one contained a plasmid with an insert of 5 kb that hybridized strongly with the probe. A map of restriction endonuclease sites was constructed and suitable fragments were cloned and sequenced (Fig. 1A). The nucleotide sequence of *ATPδ* and flanking regions, together with the deduced amino-acid sequence, is illustrated in Fig. 2. An open reading frame of 158 codons is present that has 70% nucleotide, and 72% amino-acid, identity to the *ATPδ* gene from *S. cerevisiae* (Fig. 3; Giraud and Velours 1994). Of note is the absence of a codon in the middle of the gene which in *S. cerevisiae* specifies asparagine. The amino-terminus of the deduced protein contains basic and hydrophobic amino acids that are elements of a mitochondrial targeting signal. In *S. cerevisiae* the mature δ -subunit begins with alanine which is preceded by codons specifying a maturation site (R-X-Y). Likewise, these sequence elements are conserved in the δ -subunit of *K. lactis* suggesting that a similar mature protein is likely to be produced (Fig. 3).

Disruption of *ATPδ*

To disrupt the *ATPδ* gene, a plasmid was constructed containing the *URA3* gene of *S. cerevisiae* inserted at a *Bgl*II site within the coding region (Fig. 1A). A fragment containing the disrupted *ATPδ* gene, produced by digestion with *MscI*-*Bsp*HI, was then used to transform the *K. lactis* strain PM6-7A. Transformants were selected on uracil-free




Fig. 1 A Physical map of a 5-kb fragment of *K. lactis* genomic DNA containing the *ATPδ* gene. The position of the gene and the direction of transcription is indicated by the black arrow. For gene disruption, a 1.1-kb DNA segment, containing the *S. cerevisiae* *URA3* gene, has been inserted at the *Bgl*II site. B Southern-blot analysis of gene disruption. Genomic DNA from the parental strain, PM6-7A, lane 1, and the transformant, lane 2, have been digested with *MscI* and *Bsp*HI and hybridized with [³²P]-labelled *ATPδ*

medium and examined for replacement of the resident *ATPδ* gene by gel electrophoresis and Southern-blot analysis. As illustrated in Fig. 1B, a *Ura*⁺ transformant has a DNA fragment, hybridizing to the *ATPδ* probe, that has increased in size by 1.1 kb due to the insertion of *URA3*.

Strains containing a disrupted *ATPδ* do not grow on glycerol and show a highly sectorized colony morphology on glucose (data not shown). Sectoring is probably a consequence of low viability. Micro-manipulation of individual cells shows that only 3% of cells in a liquid culture at the end of exponential growth form colonies whereas 68% of cells in a culture of the parent, PM6-7A, form colonies. Viable cells still contain mtDNA since transformation with a plasmid containing the *ATPδ* gene from *S. cerevisiae* leads to the restoration of growth on glycerol; however, transformants do not show resistance to EB. Complementation by the *S. cerevisiae* gene confirms that *ATPδ* of *K. lactis* encodes the F₁ δ -subunit. The reciprocal experiment was not attempted because *S. cerevisiae* strains with a dis-

Fig. 2 Nucleotide sequence of the *K. lactis* *ATPδ* gene and deduced amino-acid sequence. Key restriction endonuclease sites used in gene disruption have been included

AAAATGAAAAGAATGGGATTTTAGTTTTAATTACAGCTTTACGATATTGTGAATAAGGTTTACTAGAACTGTTCTTG	GATTTTGTG	-235
	<i>EcoRI</i>	-157
TTGGACTACTGTGTTCCTTCTTAGACTTTTGTATTTTGATCTTTGTTTAGTGTATTAGATTTTGTTCCTCAATTGAAT		-79
TCATTACAGTCTGTTTAAATGCTTCTGTCCAGAAATCGAAATCATAGAACACATTACCAGAAAATAATCACCTATA		-1
<i>MscI</i>		
ATGTTCCGTTTATCTGCTGCTAGAACTTTGGCCAAAATCTGTCAACACCGTGTGTTGCTAAGCGTACTTATGCTGAAGCT		78
M F R L S A A R T L A K S V N T V V A K R T Y A E A		26
GCCGATGGTGTCTTGAAGTTGCAATTTGCTTTGCCACATCAAACCTTGTTCCTGGTACTCCAGTTACCCAAGTCAAC		156
A D G A L K L Q F A L P H Q T L F S G T P V T Q V N		52
TTGCCAGCTAAGTCTGGTCAAATCGGTATCTTGGCTAACCACGTTCCAACCTGTTGAACAATTGGTTCAGGTGTTGTT		234
L P A K S G Q I G I L A N H V P T V E Q L V P G V V		78
GAAGTCTTGGAAAGGTTCTTCTAAGAAGTCTTTGTTTCCGGTGGTTTCGCTACTGTCCAACCAGACTCTACCTTGGCT		312
E V L E G S S K K F F V S G G F A T V Q P D S T L A		104
<i>BglII</i>		
ATCACTTCCGTTGAAGCATTCCCATTTAGAGTCTTTCTCTCCAGAAAACGTTAGATCTTGTGGCTGAAGCTCAAAAG		390
I T S V E A F P L E S F S P E N V R S L L A E A Q K		130
AACGCTCTCTCTGCTGACGAAGTTGCTGCTGCCGAAGCTGCTATCCAACCTGAAGTTTGGAAAGCTTTGCAAGCCGCT		468
N V S S A D E V A A A E A A I Q L E V L E A L Q A A		156
TTGAAATAATTGAGATGAACGAAGAATTACAAATGGACCAAGGTATAATACACTTGATACGGATACTCTACTTGAAG		546
L K *		158
CATTGTTTTCCCGGTTGAATGTAAACCTGTCTCGGCGTTTGTATTATCCTTCGCTGCTTTTTTTTTCATTATTCTAAT		624
<i>BspHI</i>		
TCCTATTATTCCATTGGCACAAGCTTACACAACACACGACGAATACATTTCATGATTAGCCTAAAGACCCTTTTTT		702
GTGCTCCGGTCATTGTCTTCTTTCCATACATACTGTAGTGTGAACGT		753

Kdelta	1	MFRLSAARTLAKSVNTVVAKRITYAEADG--ALKLQFALPHQTLFSGTPVTQVNLPAKSG
scdelta	1	MLRSIIKRSASRSLN-FVAKRSYAEAAASGLKLQFALPHETLYSGSEVTQVNLPAKSG
Kdelta	59	QIGILANHVPTVEQLVPGVVEVLEGS-SKKFFVSGGFATVQPDSTLAITSVEAFPLESFS
scdelta	60	RIGYLANHVPTVEQLVPGVVEVMEGSNSKKFFISGGFATVQPDSTLCVTAIEAFPLESFS
Kdelta	118	PENVRSLLEAQKNVSSADEVAEEAAIQLEVLLEALQAALK
scdelta	120	QENIKNLLAEAKKNVSSDAREAAEEAAIQVEVLENLQSVLK

Fig. 3 Alignment of *F₁* δ -subunit sequences from *K. lactis* (158 amino acids) and *S. cerevisiae* (160 amino acids). Black shading indicates identity and grey represents conserved amino acids. The two proteins have 114 identical amino acids. An arrow indicates the cleavage site of the *S. cerevisiae* precursor protein

rupted *ATPδ* do not possess functional mtDNA due to their total conversion to petites (Giraud and Velours 1994). In view of this observation with *S. cerevisiae* it is likely that low viability and colony sectoring on disruption of *ATPδ* in *K. lactis* is due to instability of mtDNA leading to the formation of inviable petite mutants.

A similar colony sectoring phenotype has been found on disruption of the *MRF1* gene encoding a mitochondrial peptide release factor in *K. lactis* (Pel et al. 1996). Low viability of *MRF1* disruptants is viewed as arising from the formation of cytoplasmic petites that do not survive. In *S. cerevisiae*, inactivation of *MRF1* leads to excessive production of petite mutants (Pel et al. 1992), which accords with the observation that interference in mitochondrial translation results in the de-stabilization of mtDNA (Myers et al. 1995).

Disruption of *ATPδ* in *K. lactis* does not alter the *Mgi*⁺ phenotype since strains do not grow in the presence of 16 µg/ml of EB (Fig. 4). In this respect, disruption of *ATPδ* produces the same *Mgi*⁺ phenotype as null mutants of *MGII*, 2 and 5 encoding the α -, β - and γ -subunits of *F₁* (Chen and Clark-Walker 1995, 1996). However a notable difference between disruptants of *ATPδ* and the three other loci is observed in the presence of a *mgi* allele.

Disruption of *ATPδ* in different *mgi* strains

Previous studies have shown that disruption of any of the three *MGI* genes in *mgi* mutants abolishes the *Mgi*⁻ phenotype. Such strains are no longer resistant to EB at 16 µg/ml, nor are petite mutants produced. However disruption of *ATPδ* in *mgi* mutants is not so simple. Strains containing different *mgi* alleles (Table 1) respond differently on disruption of *ATPδ*. Transformants of the various *mgi* mutant strains were checked for replacement of the resident *ATPδ* by Southern blotting and hybridization (data not shown) and examined for growth on glycerol and their




Fig. 4 Growth of *K. lactis* strains on glucose (GYP), glycerol (GlyYP) and response to ethidium bromide at 16 µg/ml. Cells were grown to stationary phase, diluted to approximately 10^4 cells/ml and 10-µl aliquots applied to plates which were incubated at 28°C for 5 days (GYP and GlyYP) and 7 days for GYP plus EB. Genotypes of strains are listed in Table 1

response to EB (Fig. 4). All *mgi* strains, which were chosen because of their Gly⁺ phenotype, fail to grow on glycerol after disruption of *ATPδ*, in accord with the requirement of this subunit for oxidative phosphorylation. However, growth does occur on EB with *ATPδ*-disrupted strains containing *mgi2-1*, *mex1-1* and to a lesser extent with *mgi1-9* and *mgi5-2*, whereas a Mgi⁻ phenotype is not present in disrupted strains with *mgi1-1* and *mgi2-6* mutations.

While these results indicate that the Atpδ subunit is not absolutely required for a Mgi⁻ phenotype, there is a clear need for this protein in some *mgi* mutants. In this respect it is noteworthy that assembly of a core F₁ particle, com-

posed of α-, β- and γ-subunits, can still occur in *Escherichia coli* in the absence of the ε-subunit (equivalent to δ in yeasts) although energy linked functions are abolished (Klionsky and Simoni 1985). Hence, by analogy with *E. coli*, a core F₁ particle, lacking the δ-subunit, may form in the mitochondria of some *mgi* mutants and be sufficient for suppression of ρ⁰ lethality. Alternatively, in strains with different *mgi* alleles, assembly or stabilization of a core F₁ particle could be dependent on the presence of Atpδ.

Acknowledgements We thank J. Velours for plasmid pMFG4-*ATPδ* containing *S. cerevisiae ATPδ*, and Lijun Ouyang for skilled technical assistance.

References

- Abrahams JP, Leslie AGW, Lutter R, Walker JE (1994) Structure at 2.8 Å resolution of F₁-ATPase from bovine heart mitochondria. *Nature* 370:621–628
- Aggeler R, Capaldi RA (1996) Nucleotide-dependent movement of the ε-subunit between αβ-subunits in the *Escherichia coli* F₁F₀ type ATPase. *J Biol Chem* 271:13888–13891
- Attardi G, Schatz G (1988) Biogenesis of mitochondria. *Annu Rev Cell Biol* 4:289–333
- Boyer PD (1993) The binding change mechanism for ATP synthase – some probabilities and possibilities. *Biochim Biophys Acta* 1140:215–250
- Bulder CGE (1964 a) Induction of petite mutation and inhibition of synthesis of respiratory enzymes in various yeasts. *Ant van Leeuwenhoek* 30:1–9
- Bulder CGE (1964 b) Lethality of the petite mutation in petite-negative yeasts. *Ant van Leeuwenhoek* 30:442–445
- Chen XJ (1996) Low- and high-copy number shuttle vectors for replication in the budding yeast *Kluyveromyces lactis*. *Gene* 172:131–136
- Chen XJ, Clark-Walker GD (1993) Mutations in *MGI* genes convert *Kluyveromyces lactis* into a petite-positive yeast. *Genetics* 133:517–525
- Chen XJ, Clark-Walker GD (1995) Specific mutations in α- and γ-subunits of F₁-ATPase affect mitochondrial genome integrity in the petite-negative yeast *Kluyveromyces lactis*. *EMBO J* 14:3277–3286
- Chen XJ, Clark-Walker GD (1996) The mitochondrial genome integrity gene *MGI1* of *Kluyveromyces lactis* encodes the β-subunit of F₁-ATPase. *Genetics* 144:1445–1454
- Chen XJ, Wesolowski-Louvel M, Fukuhara H (1992) Glucose transport in the yeast *Kluyveromyces lactis*. II. Transcriptional regulation of the glucose transporter gene *RAG1*. *Mol Gen Genet* 233:97–105
- Clark-Walker GD, Chen XJ (1996) A vital function for mitochondrial DNA in the petite-negative yeast *Kluyveromyces lactis*. *Mol Gen Genet* 252:746–750
- Clark-Walker GD, McArthur CR, Daley DJ (1981) Does mitochondrial DNA length influence the frequency of spontaneous petite mutations in yeasts? *Curr Genet* 4:7–12
- Cox GB, Devenish RJ, Gibson F, Howitt SM, Nagley P (1992) The structure and assembly of ATP synthase. In: Ernster L (ed) *Molecular mechanisms in bioenergetics*. Elsevier Science Publication BV, pp 283–315
- De Deken RH (1966) The Crabtree effect and its relation to the petite mutation. *J Gen Microbiol* 44:157–165
- Deckers-Hebestreit G, Altendorf K (1996) The F₀ F₁-type ATP synthases of bacteria: structure and function of the F₀ complex. *Annu Rev Microbiol* 50:791–824
- Duncan TM, Bulygin VV, Zhou Y, Hutcheon ML, Cross RL (1995) Rotation of subunits during catalysis by *Escherichia coli* F₁-ATPase. *Proc Natl Acad Sci USA* 92:10964–10968

- Ephrussi B (1953) Nucleocytoplasmic relations in microorganisms. Clarendon Press, Oxford
- Futai M, Noumi T, Maeda M (1989) ATP synthase (H^+ -ATPase): results by combined biochemical and molecular biological approaches. *Annu Rev Biochem* 58:111–136
- Giraud M-F, Velours J (1994) ATP synthase of yeast mitochondria. Isolation of the F_1 δ -subunit sequence and disruption of the structural gene. *Eur J Biochem* 222:851–859
- Grivell LA (1989) Nucleo-mitochondrial interactions in yeast mitochondrial biogenesis. *Eur J Biochem* 182:477–493
- Hatefi Y (1993) ATP synthesis in mitochondria. *Eur J Biochem* 218:759–769
- Heritage J, Whittaker PA (1977) Isolation of metabolically active petite mutants of *Kluyveromyces lactis* a petite-negative yeast. *Mol Gen Genet* 156:93–98
- Klionsky DJ, Simoni RD (1985) Assembly of a functional F_1 of the proton-translocating ATPase of *Escherichia coli*. *J Biol Chem* 260:11200–11206
- Mullis KB, Faloona FA (1987) Specific synthesis of DNA in vitro via a polymerase-catalysed chain reaction. *Methods Enzymol* 155:335–350
- Myers AM, Pape LK, Tzagoloff A (1985) Mitochondrial protein synthesis is required for maintenance of intact mitochondrial genomes in *Saccharomyces cerevisiae*. *EMBO J* 4:2087–2092
- Noji H, Yasuda R, Yoshida M, Kinoshita Jr K (1997) Direct observation of the rotation of F_1 -ATPase. *Nature* 386:299–302
- Pel HJ, Maat C, Rep M, Grivell LA (1992) The yeast nuclear gene *MRF1* encodes a mitochondrial peptide chain release factor and cures several mitochondrial RNA splicing defects. *Nucleic Acids Res* 20:6339–6346
- Pel HJ, Rozenfeld S, Bolotin-Fukuhara M (1996) The nuclear *Kluyveromyces lactis* *MRF1* gene encodes a mitochondrial class-I peptide chain release factor that is important for cell viability. *Curr Genet* 30:19–28
- Poyton RO, McEwen JE (1996) Crosstalk between nuclear and mitochondrial genomes. *Annu Rev Biochem* 65:563–607
- Rothstein RJ (1983) A one-step gene disruption in yeast. *Methods Enzymol* 101:202–210
- Sabbert D, Engelbrecht S, Junge W (1996) Inter-subunit rotation in active F_1 -ATPase. *Nature* 381:623–625
- Sanger FS, Nicklen S, Coulson AR (1977) DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci USA* 74:5463–5467
- Senior AE (1990) The proton-translocating ATPase of *Escherichia coli*. *Annu Rev Biophys Biophys Chem* 19:7–41
- Slonimski P, Perrodin G, Croft JH (1968) Ethidium bromide-induced mutation of yeast mitochondria: complete transformation of cells into respiratory deficient non-chromosomal 'petites'. *Biochem Biophys Res Commun* 30:232–239
- Walker JE, Collinson IR (1994) The role of the stalk in the coupling mechanism of $F_1 F_0$ -ATPases. *FEBS Lett* 346:39–43
- Wesolowski-Louvel M, Tanguy-Rougeau C, Fukuhara H (1986) A nuclear gene required for the expression of the linear DNA-associated killer system in the yeast *Kluyveromyces lactis*. *Yeast* 4:71–81

Enolase and Glycolytic Flux Play a Role in the Regulation of the Glucose Permease Gene *RAG1* of *Kluyveromyces lactis*

Marc Lemaire¹ and Micheline Wésolowski-Louvel

UMR 5122 Microbiologie et Génétique, Université Claude Bernard Lyon 1, 69622 Villeurbanne, France

Manuscript received April 13, 2004

Accepted for publication June 3, 2004

ABSTRACT

We isolated a mutant, *rag17*, which is impaired in glucose induction of expression of the major glucose transporter gene *RAG1*. The *RAG17* gene encodes a protein 87% identical to *S. cerevisiae* enolases (Eno1 and Eno2). The *Kleno* null mutant showed no detectable enolase enzymatic activity and has severe growth defects on glucose and gluconeogenic carbon sources, indicating that *K. lactis* has a single enolase gene. In addition to *RAG1*, the transcription of several glycolytic genes was also strongly reduced in the Δ *Kleno* mutant. Moreover, the defect in *RAG1* expression was observed in other mutants of the glycolytic pathway (hexokinase and phosphoglycerate kinase). Therefore, it seems that the enolase and a functional glycolytic flux are necessary for induction of expression of the *Rag1* glucose permease in *K. lactis*.

IN most strains of *Kluyveromyces lactis*, the glucose uptake system relies on two nonredundant glucose transporters: a low-affinity permease encoded by *RAG1* (WÉSOLOWSKI-LOUVEL *et al.* 1992a) and a high-affinity permease encoded by *HGT1* (BILLARD *et al.* 1996). *HGT1* is constitutively expressed (BILLARD *et al.* 1996); expression of *RAG1* is activated in the presence of high concentrations of glucose (CHEN *et al.* 1992; WÉSOLOWSKI-LOUVEL *et al.* 1992a). The *Rag1* permease is necessary for supporting fermentative growth, which requires a high flow of substrate. In the absence of *Rag1*, the cell becomes respiration dependent for growth on high-glucose media. Thus, *rag1* cells have the *Rag*[−] phenotype: they cannot grow on 5% glucose in the presence of antimycin A, which blocks respiration (GOFFRINI *et al.* 1989; WÉSOLOWSKI-LOUVEL *et al.* 1992b).

To date, studies of *Rag*[−] mutants have identified three key components that are involved in the positive regulation of *RAG1* expression: the glucose sensor *Rag4* (BETINA *et al.* 2001), hexokinase *Rag5* (PRIOR *et al.* 1993), and casein kinase I *Rag8* (BLAISONNEAU *et al.* 1997).

In this report we present the characterization of another gene in *K. lactis* implicated in *RAG1* regulation: *RAG17* (*KIENO*) coding for enolase.

MATERIALS AND METHODS

Yeast strains and growth conditions: Yeast strains are described in Table 1. Yeast cells were grown at 28° in a complete YP medium containing 1% Bacto yeast extract, 1% Bactopeptone (Difco, Detroit), supplemented with either 2% glucose (YPG) or a specified carbon source. Minimal medium containing 0.7% yeast nitrogen base without amino acids (Difco) and 2% glucose was supplemented with auxotrophic requirements. The *Rag* phenotype was tested on 5% complete glucose medium supplemented with 5 μ M antimycin A. For G418 medium, YPG plates were supplemented with geneticin (200 μ g/ml; Life Technologies). 5-FOA medium was prepared according to BOERKE *et al.* (1984). The media for plates were solidified by the addition of 2% Bactoagar (Difco). *Escherichia coli* XL1-blue was used as a cloning host and DNA propagation and was grown in LB medium.

Genetics methods have been described previously (WÉSOLOWSKI *et al.* 1982; GOFFRINI *et al.* 1989).

Yeast transformation: Replicative transformation of *K. lactis* was performed by electroporation. For integrative transformation of *K. lactis*, the procedure described by DOHMEN *et al.* (1991) was followed. Replicative and integrative transformations of *Saccharomyces cerevisiae* were standard.

Construction of deletion strains: One-step gene deletions using *kanMX4* or *HIS3* selection markers bearing PCR-generated long flanking homology (LFH-PCR; WACH 1996) was used to construct the mutant strains MWK5, MLK43, MLY702, and MLY703 (Table 1). The primers used for the *KIENO* LFH-PCR synthesis were P5' *KIENO* (5'-CACGTTTCAATCCAGGC ACC-3'), P5' *L KIENO* (5'-CCGTCGACCTGCAGCGTACGTGG CATGTTTTTTT-3'), P3' *L KIENO* (5'-GCTCGAATTCATC GATGATATTTGACTGTCCACCAAC-3'), and P3' *KIENO* (5'-AGC GAAGATAGCGTTGGAAACC-3'). The primers used for the *KIPGK* LFH-PCR synthesis were P5' *KIPGK* (5'-ACGATCTCGTCTAG TGGAAACC-3'), P5' *L KIPGK* (5'-GGGATCCGTCGACCTGC AGCGTACGATTTTTTATTAATTCITGATCG-3'), P3' *L KIPGK* (5'-CGAGCTCGAATTCATCGATGATATAAATGTAGGATC CATCATCCC-3'), and P3' *KIPGK* (5'-TACGATGAACCACTGCA CAAG-3'). The primers used for the *SENO2* LFH-PCR synthesis

Sequence data from this article have been deposited with the EMBL Data Library under accession no. AJ586240.

¹Corresponding author: Unité Microbiologie et Génétique UMR 5122 CNRS/UCBL/INSA, Université Claude Bernard, Bâtiment Lwoff, 43, Blvd. du 11 Novembre 1918, 69622 Villeurbanne Cédex, France. E-mail: mlemaire@univ-lyon1.fr

were P5' *ScENO2* (5'-ATCCTACTCTTCCGCTTGCCATCC-3'), P5' *L. ScENO2* (5'-GGGGATCCGTCGACCTGCAGCGTACGCA TTATTATTGTAT GTTATAGTA-3'), P3' *L. ScENO2* (5'-AAACGA GCTCGAATTCATCGATGATATAAAGTGCTTTTAACTAAGA ATT-3'), and P3' *ScENO2* (5'-GTACTCATAGAGGTAGGCTAGA CC-3'). For all the P5' and P3' primers, the *KanMX4* or *HIS3* MX6 regions are in boldface type. All correct integrations were verified by Southern blot or PCR.

In *S. cerevisiae*, the *ScENO2* gene was first disrupted in the diploid strain MLY701 by using a *kanMX4* or a *HIS3MX6* disruption cassette, yielding MLY702 and MLY703 strains, respectively (Table 1). After meiotic analysis of these two diploid strains, the Δ *Sceno2* haploid strains MLY704 (Δ *Sceno2::kanMX4*) and MLY708 (Δ *Sceno2::HIS3*) were obtained. The double null mutant strain Δ *Sceno1* Δ *Sceno2* was then constructed by crossing the MLY708 strain with the Δ *Sceno1* strain Y07286 (Table 1), yielding the MLY713 diploid.

In *K. lactis*, the *KIENO* and *KIPGK* genes were disrupted in the MW270-7B strain using the corresponding *kanMX4* disruption cassettes, yielding MWK3 and MLK43 strains, respectively.

Plasmid constructions: The *ENO2* gene of *S. cerevisiae* was first PCR amplified from MLY701 genomic DNA by using the Pfx polymerase (Invitrogen, San Diego) and the P5' *ScENO2*/P3' *ScENO2* primers. After phosphorylation with T4 polynucleotide kinase (New England Biolabs, Beverly, MA), the *ScENO2* PCR product was cloned into the *CEN-URA3* vector pRS416 (STKORSKI and HUFER 1989) linearized with *SmaI*, yielding pML180. This plasmid could complement the slow growth of the Δ *Sceno2* mutant (MLY708, Table 1) and thus contained a functional copy of *ScENO2*. The 2.9-kb *XhoI*-*NofI* fragment of pML180, containing *ScENO2*, was then cloned into the *CEN TRP1* vector pRS414 (STKORSKI and HUFER 1989) digested with *XhoI* and *NofI*, yielding pML190. The *ScENO2* was also introduced into a *K. lactis* centromeric vector by cloning the *XbaI*-*XhoI* 2.9-kb fragment of pML180 between the *XbaI* and *SalI* sites of the *CEN URA3* pCXJ18 vector (CHEN 1996), yielding pML187.

The *K. lactis* *KIENO* gene was cloned into a *S. cerevisiae* vector by subcloning the 5.5-kb *BamHI* fragment of pMW1 (see below) into the *BamHI* site of pRS414, yielding pML183.

Cloning and sequencing of *RAG17/KIENO* gene: The *RAG17* gene was cloned by *in vivo* complementation of *rag17* mutation (strain MWK2) with a *K. lactis* genomic library made in the KCP491 vector (PRIOR *et al.* 1993). Of 5000 *Ura*⁺ transformants, 4 were found to be Rag⁺. The complementing plasmids extracted from these four transformants and amplified in *E. coli* showed that three of them carried the same plasmid, pMW1 containing a 7.5-kb insert (Figure 1), and one a plasmid with a 10-kb insert overlapping with a pMW1 insert.

A 2856-bp fragment containing the entire *RAG17/KIENO* gene was sequenced on both strands.

Preparation of yeast RNA and probes: Total RNA was extracted from cells grown to an OD₆₀₀ of 2–3. Poly(A⁺)-enriched mRNA was obtained using mRNA Separator (CLONTECH, Palo Alto, CA). Several gene probes used were restriction fragments: *RAG6* probe was a 1.3-kb *EcoRI*-*SalI* fragment (BIANCHI *et al.* 1996); *HGT1* probe was a 1.75-kb *EcoRI*-*HindIII* fragment containing the *HGT1* gene (BILLARD *et al.* 1996). Other probes were obtained by PCR amplification using either *K. lactis* genomic DNA or the cloned genes as template. The oligonucleotides used were: 0487 (5'-GGGGTCGTAGAAATTGGT-3') and 0369 (5'-GACGTAACCGTAGTAGAAG-3') for *RAG1*; *RAG2*-up (5'-TGTACGTTGATCGGTACCAACG-3') and *RAG2*-down (5'-CAAG ATAGAACCACTAGAGTA-3') for *RAG2*; p2E1/9 (5'-GCCATCT GTGCAGCATCAA-3') and p2E1/5 (5'-GGGAAGAAGATCGAG TAGTG-3') for *RAG4*; 470 (5'-GTGCCAGCTAATTTGATGGA-3') and 471 (5'-ACCAGCCACCAATTGGATTG-3') for *RAG5*;

PENO1 (5'-ACTGGCTGTCTGACTAGC-3') and *EndoY* (5'-GTC TTAGCACCGGCCAAGTC-3') for *KIENO*; *SCK1*-up (5'-GAACAC CAACATGTTGCTACTC-3') and *SCK1*-down (5'-GACAACG AACCCAGTATCTTCGC-3') for *SCK1*; *GCR1*-up (5'-CACCAG TAACATGATACGGTCC-3') and *GCR1*-down (5'-GACCACCAT CAGATATACTGTTGCC-3') for *KIGCR1*; *GCR2*-B (5'-TCAGC GATTTCAACAGATAT-3') and *GCR2*-D (5'-CTCATTGATCTGT TCCATAG-3') for *KIGCR2*. In all cases, specific probes of *K. lactis* actin gene (*KIAC1*), *KhRNA 18S* gene (*18S*), or *KIAAC* gene were used in parallel as quantitative references. In the case of *KIAC1*, the probe was a 900-bp *EcoRI*-*HindIII* fragment. *18S* and *KIAAC* probes were amplified by PCR using the following oligonucleotides: PKI-18S f (5'-ATCCTGCCAGTAGTCATATGC-3') and PKI-18S r (5'-CCACAAGGAGTACAGGTTAGC-3') for *rRNA 18S*; P5' *KIAAC* (5'-AGATGAAATGATCAAGCAAGG-3') and P3' *KIAAC* (5'-CGTACATGGAGATAACCCGG-3') for *KIAAC*.

Northern blot hybridization was quantified by scanning with a Cyclone Phosphorimager (Packard, Meriden, CT).

Preparation of cell-free extract and enzyme assays: Whole-yeast-cell extracts were prepared by glass-bead disruption of cells isolated from log-phase cultures grown in YP medium containing either 2% glucose or 2% glycerol. Enolase activity was assayed (CLIFTON *et al.* 1978) and normalized to protein concentration determined by the Bradford protein assay (Bio-Rad, Richmond, CA).

RESULTS

Isolation of the *KIENO* gene and deduced amino acid sequence of its product: In *K. lactis*, gene replacement by homologous recombination can be accomplished, but at lower frequencies compared to *S. cerevisiae*. Usually, the gene disruption cassette recombines at ectopic sites in the genome. While attempting to construct a *rag4::URA3* gene disruption (BETINA *et al.* 2001), we identified a Rag⁻ mutant that is not allelic to *rag4*, although *RAG1* transcription is highly reduced in this mutant. The mutation is also not allelic to *rag5* and *rag8* mutations, both of which affect genes that positively regulate the transcription of the *RAG1* gene (CHEN *et al.* 1992). We named the mutation *rag17-1* (strain MWK1 in Table 1). The precise position of the *URA3* insertion in the gene is not known.

RAG17 was isolated from a *CEN*-based *K. lactis* plasmid library by complementation of the Rag⁻ phenotype of the *rag17-1* mutation (MATERIALS AND METHODS). The partial nucleotide sequence of the DNA fragment in the complementing plasmid (Figure 1 and MATERIALS AND METHODS) revealed the presence of three ORFs: one encodes a protein 40% identical to the Fmo of *S. cerevisiae*, a flavin-containing monooxygenase involved in protein folding (SUH *et al.* 1999); another encodes a protein highly similar to the enolases of *S. cerevisiae*, Enol and Eno2 (HOLLAND *et al.* 1981); the third encodes a protein 28% identical to Rax1 of *S. cerevisiae*, a protein implicated in bud site selection (CHEN *et al.* 2000). Because of the Rag⁻ phenotype of the mutation, the best candidate for *RAG17* was the ORF that encodes the glycolytic enzyme enolase (*KIENO*). The predicted protein (437 amino acids) is 88 and 87% identical to *ScEno2*

TABLE 1
Yeast strains

Strain	Relevant genotype	Reference
<i>K. lactis</i>		
MW270-7B	<i>MATa ura1-1 met1-1 leu2</i>	BILLARD <i>et al.</i> (1996)
PM4-4B	<i>MATx ura1-1 ade2-1</i>	GOFFRNI <i>et al.</i> (1991)
PM6-7A	<i>MATa ura1-1 ade1-600</i>	CHEN <i>et al.</i> (1992)
MWK1	Isogenic to MW270-7B except <i>Kleno::URA3 (rag17-1)</i>	This work
MWK2	Isogenic to MW270-7B except <i>Kleno::ura3</i>	This work
MWK3	Isogenic to MW270-7B except <i>KlenoΔ1::kanMX4</i>	This work
MW352-2D	<i>MATa met1-1 ade2-1 Kleno::URA3</i>	This work
MLK43	Isogenic to MW270-7B except <i>Klpgk Δ1::kanMX4</i>	This work
MWK11	Isogenic to MW270-7B except <i>rag5Δ2::URA3 (ΔKlthsk)</i>	This work
MWK12	Isogenic to PM4-4B except <i>rag2Δ1::URA3 (ΔKlpgi)</i>	GOFFRNI <i>et al.</i> (1991)
MWK13	Isogenic to PM6-7A except <i>rag6Δ1::URA3 (ΔKlpgi)</i>	BIANCHI <i>et al.</i> (1996)
<i>S. cerevisiae</i>		
Y07286	<i>MATa ura3 Δ0 met15 Δ0 leu2 Δ0 his3Δ1 YGR254w^a::kanMX4</i>	Euroscarf
BM64-1A	<i>MATa ura3-1 trp1Δ2 leu2-3,112 his3-11 ade2-1 can1-100</i>	Euroscarf
BM64-1B	<i>MATa ura3-1 trp1Δ2 leu2-3,112 his3-11 ade2-1 can1-100</i>	Euroscarf
MLY701	Diploid issued from BM64-1A × BM64-1B cross	This work
MLY702	Isogenic to MLY701 except <i>ENO2/eno2Δ1::kanMX4</i>	This work
MLY703	Isogenic to MLY701 except <i>ENO2/eno2Δ2::HIS3</i>	This work
MLY704	Meiotic segregant of MLY702 diploid: <i>MATa ura3-1 trp1Δ2 leu2-3,112 his3-11 ade2-1 can1-100 eno2Δ1::kanMX4</i>	This work
MLY708	Meiotic segregant of MLY703 diploid: <i>MATa ura3-1 trp1Δ2 leu2-3,112 his3-11 ade2-1 can1-100 eno2Δ2::HIS3</i>	This work
MLY713	Diploid issued from Y07286 × MLY708 cross	This work
MLY714	MLY713 diploid transformed with pML180 (<i>ScENO2/pRS416</i>)	This work
MLY719	Meiotic segregant of MLY714 diploid: <i>MATa ura3-1 trp1Δ2 leu2-3,112 his3-11 ade2-1 YGR254w^a::kanMX4 eno2Δ2::HIS3 + pML180 (ScENO2/pRS416)</i>	This work

^a *YGR254w*, *ScENO1*.

and ScEno1 of *S. cerevisiae*, respectively, and 64% identical to human α -enolase (Swiss-Prot accession P06733).

A PCR-based gene deletion cassette for *RAG17/KIENO* (Figure 1; Table 1; see MATERIALS AND METHODS

for details) was used to disrupt the gene in the MW270-7B strain. Southern blot analysis confirmed the *KIENO* disruption in some G418^r transformants, such as MWK3 (data not shown). We also demonstrated that the *KIENO*




FIGURE 1.—(A) Restriction map of recombinant plasmid pMW1 and localization of the three ORFs (large arrows) identified by nucleotide sequences. Small arrows indicate the direction of DNA sequencing. (B) Restriction map of the *RAG17/KIENO* locus. (C) Disruption cassette of the *KIENO* gene with *kanMX4* marker (open box) to construct the MWK3 strain (Table 1). The solid boxes indicate genomic sequences (see MATERIALS AND METHODS).



FIGURE 2.—Growth phenotype of the *Kleno* null mutant strain. The wild-type strain MW270-7B (*KIENO*), and the isogenic mutant strain MWK3 ($\Delta Kleno$) were streaked onto single colonies on 2% glucose, 2% glycerol, and 2% ethanol minimal plates. The photographs were taken after 3 days of incubation at 28°.

locus is modified in the original *rag17-1* mutant (MWK1), suggesting an ectopic integration of the *rag4::URA3* gene disruption cassette in the *KIENO* gene. Like the *rag17-1* mutation, the *Kleno* null mutation leads to a *Rag*[−] phenotype. The allelism of *Kleno* with *rag17-1* was confirmed by the absence of complementation of the *Kleno* null mutation (strain MWK3) by the *rag17-1* mutation (MW352-2D) in a diploid constructed by crossing these two mutants. Thus, the cloned *KIENO* gene indeed corresponds to the *RAG17* locus.

Growth phenotype of the *Kleno* mutant: Growth of the *Kleno* null mutant is severely reduced on media containing either glucose or glycerol as the sole carbon

source (Figure 2). In addition, the mutant strain is unable to grow on ethanol as carbon source. It is noteworthy that the *Kleno* mutant can grow slowly on complete glucose medium: it exhibits a 4.5-fold increase of doubling time as compared to wild-type cells (450 min vs. 110 min; data not shown). Better growth of mutant cells on complete medium is probably supported by other carbon sources (e.g., amino acids) in this medium.

Enolase activity of *Kleno* mutant and *KIENO* expression: No detectable enolase activity is present in the $\Delta Kleno$ mutant, regardless of the substrate used (glycerol or glucose; Figure 3A). This strongly suggests that *KIENO* is the single enolase-encoding gene in *K. lactis*. Presence of the *KIENO* sequence in a single copy and absence of other related sequences in the genome was confirmed by low-stringency Southern blotting (data not shown). In contrast, in *S. cerevisiae*, which possesses two enolase genes, enolase activity is still detectable in $\Delta Sceno1$ or $\Delta Sceno2$ single mutants grown on glucose or on glycerol. It was not possible to assay enolase activity in the double-mutant $\Delta Sceno1\Delta Sceno2$ since it is inviable on glucose or glycerol media (see below).

KIENO transcription was examined by a Northern blot analysis. The level of the *KIENO* transcript is slightly higher (twofold) when the cells are grown on 2% glucose (Figure 3B) than when they are cultivated on 2% glycerol. This result is consistent with the increased level of enolase activity detected in glucose-grown cells as compared to glycerol-grown cells (Figure 3A). In the null mutant, no transcript could be detected whatever the carbon source used. These results demonstrate that *K. lactis* possesses the single gene *KIENO* coding for an enolase and expressed under glycolytic as well as neoglucogenic conditions.

The *KIENO* gene complements the $\Delta Sceno1 \Delta Sceno2$ mutations of *S. cerevisiae*: The nucleotide sequences of the two enolase-encoding genes of *S. cerevisiae*, *ScENO1* and *ScENO2*, are >90% identical. The major difference between the two genes is in their 5' noncoding sequence. Therefore, they are expressed differently: the gluconeogenic gene *ScENO1* is constitutively expressed (COHEN *et al.* 1987); the glycolytic gene *ScENO2* is induced by glucose (MCALISTER and HOLLAND 1982; COHEN *et al.* 1986). The *Sceno1* null mutation has no phenotype (MCALISTER and HOLLAND 1982); the $\Delta Sceno2$ mutant grows more slowly on glucose than does the wild-type strain (NIEDENTHAL *et al.* 1999). Meiotic analysis of the diploid MLY713 (*ScENO2*/ $\Delta Sceno2::HIS3 *ScENO1*$




FIGURE 3.—(A) Enolase activity in wild-type and enolase mutant strains of *K. lactis* and *S. cerevisiae*. Enzyme activity was assayed as described in MATERIALS AND METHODS from cells grown on glucose or glycerol. Specific activities are expressed as micromoles of the product formed per minute and per milligram of proteins. *S. cerevisiae* strains were BM61-1A (*ScENO1 ScENO2*), MLY704 (*ScENO1 ΔSceno2*), and Y07286 ($\Delta Sceno1 ScENO2$). *K. lactis* strains were MW270-7B (*KIENO*) and MWK3 ($\Delta Kleno$). (B) Northern blot analysis of *KIENO* mRNA. Each slot was loaded with ~5–10 μg of total RNA and electrophoresed on a 1.2% agarose-formaldehyde gel. The probes used are described in MATERIALS AND METHODS. Lane 1, MW270-7B (*KIENO*) strain grown on 2% glycerol; lane 2, MW270-7B (*KIENO*) strain grown on 2% glucose; lane 3, MWK3 ($\Delta Kleno$) strain grown on 2% glycerol; lane 4, MWK3 ($\Delta Kleno$) strain grown on 2% glucose.




FIGURE 4.—Complementation of the $\Delta Sceno1\Delta Sceno2$ double mutation of *S. cerevisiae* by the *KIENO* gene. The $\Delta Sceno1\Delta Sceno2$ cells (strain MLY719, Table 1), rescued with the pML180 plasmid (*ScENO2* gene cloned in the *URA3* vector pRS416), were transformed with *TRP1* vector pRS414 (empty vector) or with the *ScENO2* and *KIENO* genes individually carried by pRS414. Transformants were grown in parallel on uracil-less, tryptophan-less, and 5-FOA medium. The plates were incubated for 3 days at 28° before the photographs were taken.

$\Delta Sceno1::kanMX4$) showed that no G418^R His⁺ spores are viable on YP medium containing either 2% glucose or 2% glycerol, but viable G418^R His⁺ spores could be obtained on YP medium containing 0.1% glucose + 2% ethanol. Under those conditions the $\Delta Sceno1\Delta Sceno2$ mutant probably obtains energy by respiration, which is derepressed at low concentrations of glucose.

To test if the *KIENO* gene can complement the growth defect of $\Delta Sceno1\Delta Sceno2$, we sporulated the MLY713 diploid (Table 1) carrying a *CEN-URA3* plasmid containing the *ScENO2* gene. A meiotic segregant of a complete tetrad, MLY719 (Ura⁺ His⁺ G418^R Trp⁻), was transformed in parallel with a *CEN-TRP1* plasmid carrying either *KIENO* or *ScENO2* (Table 1). In both cases the transformants were able to grow on 5-FOA medium, which counterselects for Ura⁺ cells (Figure 4). As a control, the same strain was transformed with the empty pRS414 plasmid and found unable to lose the plasmid carrying *ScENO2* (pML180; i.e., this strain cannot grow on 5-FOA medium). We conclude that *KIENO* restores viability to the $\Delta Sceno1\Delta Sceno2$ mutant of *S. cerevisiae*. Thus, *KIENO* is a functional homolog of the *ENO* genes of *S. cerevisiae*. The reciprocal heterologous complementation was confirmed: *ScENO2*, cloned in a *K. lactis* centromeric vector (pML187; see MATERIALS AND METHODS), complements the Rag⁻ phenotype of the *Kleno* mutant (data not shown).

KIENO is required for expression of genes encoding glucose permeases and glycolytic genes and their regulators: Northern blot analysis presented in Figure 5 shows that the disruption of *KIENO* results in a severe reduction of transcript levels of both glucose transporter genes, *RAG1* and *HGT1*. However, the transcription of *HGT1* was less affected than that of *RAG1*. The transcription of the hexokinase (*RAG5*, *KIHXX*) and pyruvate decarboxylase gene (*RAG6*, *KIPDC*) is also impaired in the mutant. In contrast, the phosphoglucose isomerase gene (*RAG2*, *KIPGI*) is not affected (Figure 5).

The reduction of the transcription of genes encoding glycolytic enzymes and glucose transporters could result from a direct effect on these genes or it could be indirect, possibly through effects on the expression of genes encoding their regulators. One of these regulators—the glucose sensor, a positive regulator of *RAG1*—is encoded by *RAG4* (BETINA *et al.* 2001). Another one, *SCK1*,

codes for a helix-loop-helix type DNA-binding transcription factor, homologous to *SGC1* of *S. cerevisiae*, which is required for full expression of glycolytic genes and of the glucose carrier gene *RAG1* (LEMAIRE *et al.* 2002). *KIGCR1* and *KIGCR2* of *K. lactis* are the orthologs of the positive regulatory genes *GCR1* and *GCR2* of *S. cerevisiae* (HAW *et al.* 2001; NEIL *et al.* 2004). The *KIGCR1* and *KIGCR2* genes, like *SCK1*, appear to positively control




FIGURE 5.—Effect of the disruption of *KIENO* on the transcription of glucose-transporter and glycolytic genes. Approximately 5–10 μ g of total RNA extracted from cells grown on 2% glucose was loaded in each slot. Strains used were MW270-7B (*KIENO*), MWK3 (*KlenoΔ1::kanMX4*), and MWK1 (carrying the original *Kleno::URA3* mutation) (Table 1). Electrophoresis conditions were as in Figure 3. The probes used to detect *RAG1*, *HGT1*, *RAG5* (*KIHXX* coding for hexokinase), *RAG2* (*KIPGI* coding for phosphoglucose isomerase), *RAG6* (*KIPDC* coding for pyruvate decarboxylase), and *18S* transcripts are described in MATERIALS AND METHODS. As already known, two mRNA are detected for *HGT1* (BILLARD *et al.* 1996). *18S* mRNA was used as an internal standard.




FIGURE 6.—Northern blot analysis of *KIGCR1*, *KIGCR2*, *SCK1*, and *RAG4* transcription in *Kleno* null mutant. Approximately 5 μ g of poly(A⁺) mRNA prepared from cells grown on 2% glucose was loaded in each slot. Strains used were MW270-7B (*Kleno*) and MWK3 (*Kleno* Δ 1::kanMX4). The probes used are described in MATERIALS AND METHODS.

the expression of glycolytic genes and *RAG1* (NEIL *et al.* 2004). The amount of each of these transcripts was decreased compared to the wild-type strain in the enolase mutant strain (Figure 6). The transcript levels of these genes were reduced approximately five- to sevenfold in the mutant relative to the reference *KIACT* and *KIAAC* transcripts (data not shown).

A robust glycolytic flux is necessary for the full expression of *RAG1*: *KIENO* is the second glycolytic gene controlling the *RAG1* regulation to be identified. The *RAG5* gene, coding for the single hexokinase in *K. lactis*, was already known to be required for the full expression of *RAG1* (PRIOR *et al.* 1993). Enolase and hexokinase are suspected to harbor both catalytic and regulatory functions (BISSON and FRAENKEL 1983; ENTIAN and FRÖHLICH 1984; PRIOR *et al.* 1993; FEO *et al.* 2000; SUBRAMANIAN and MILLER 2000), but our data suggest a more general hypothesis: the overall glycolytic flux may regulate glucose transport.

To investigate this hypothesis, we analyzed the expression of *RAG1* in several mutants that are defective for different steps of the glycolytic or fermentation pathways (Figure 7A): Δ *Klhxk* (hexokinase), Δ *Klpgi* (phosphoglucosomerase), Δ *Klpgk* (phosphoglycerate kinase), Δ *Kleno* (enolase), and Δ *Klpgc* (pyruvate decarboxylase). The results showed that *RAG1* expression is significantly reduced in Δ *Klhxk*, Δ *Klpgk*, and Δ *Kleno* mutants (Figure 7B) in which the glycolytic flux is blocked. However, the Δ *Klpgi*

mutant, which can bypass the glycolytic block through the pentose phosphate pathway (JACOBY *et al.* 1993; GONZALEZ SISO *et al.* 1996), has little or no effect. The Δ *Klpgc* mutation that blocks the first step of fermentation following glycolysis has no impact on *RAG1* transcription. These findings suggest that glycolytic flux is required for full expression of *RAG1*.

Interestingly, regulation of glucose uptake by glycolytic flux in *S. cerevisiae* has been suggested (BISSON *et al.* 1993). We tested whether the activation of *HXT1* (most closely related to *RAG1*) expression is impaired in the enolase mutant of *S. cerevisiae* grown on glucose. No effect on expression of an *HXT1-LacZ* fusion was observed in the single *eno1* or *eno2* mutants (data not shown). This negative result is not necessarily conclusive, because both single mutants retain some enolase activity (Figure 3A). Unfortunately, this experiment cannot be performed with the double-mutant *eno1 eno2* since this mutant cannot grow on glucose (see Figure 2).

A reduced growth rate does not affect *RAG1* transcription: The glycolysis block in the *Kleno* mutant leads to a severe growth defect (Figure 2). The *Klhxk* and *Klpgk* mutants, but not the *Klpgi* and *Klpgc* mutants, show a similar growth defect (GOFFRINI *et al.* 1991; PRIOR *et al.* 1993; BIANCHI *et al.* 1996; data not shown). Thus, it remained possible that the *RAG1* transcriptional defect in the *Klhxk*, *Klpgk*, and *Kleno* mutants could be due to their reduced growth rate rather than to their reduced glycolytic capability. To investigate this possibility, a wild-type strain (MW270-7B, *KIENO*) was grown in YPG with or without different growth inhibitors. We used antimycin A and potassium cyanide (KCN), which block the respiratory chain reaction, and geneticin (G418), which inhibits protein synthesis. Figure 8A shows that these compounds inhibit the growth of *K. lactis* on YPG. However, Northern blot analysis (Figure 8B) demonstrated that the *RAG1* gene is still inducible by glucose in nondividing cells in the presence of antimycin A, KCN, or G418. This demonstrates that the *RAG1* gene is induced to similar levels in dividing and nondividing cells. These results support the idea that the defect in *RAG1* transcription in the *Kleno*, *Klhxk*, and *Klpgk* mutants is caused by reduced glycolytic flux rather than by impaired growth.

DISCUSSION

We have shown that the *KIENO* gene, which encodes enolase, the glycolytic enzyme that catalyzes the reversible conversion of 2-phosphoglycerate to phosphoenolpyruvate, is required for normal regulation of expression of *RAG1*, encoding the low-affinity glucose permease in *K. lactis*. Glycolytic enzymes have been extensively studied and characterized at the structural and biochemical level. Recently, interest in glycolytic enzymes has been revived due to their implications in other biological pathways. In *S. cerevisiae*, the hexokinase PII, encoded




FIGURE 7.—Northern blot analysis of *RAG1* in mutant affected at different steps of glycolysis (A). Approximately 5–10 μ g of total RNA extracted from cells grown on 2% glucose was loaded in each slot. Electrophoresis conditions were as in Figure 3. The probes used to detect *RAG1* and *18S* transcripts are described in MATERIALS AND METHODS. (B) The strains MW270-7B, PM4-4B, and PM6-7A are the isogenic wild-type strains of the different mutants used. $\Delta Khlxk$, MWK11 strain; $\Delta Klpgl$, MWK12 strain; $\Delta Klpgh$, MWK45 strain; $\Delta Kleno$, MWK3 strain; $\Delta Klpdc$, MWK13 strain. These strains are described in Table 1. *18S* mRNA was used as an internal standard. The hybridization signals have been quantified with a phosphorimager (MATERIALS AND METHODS) and numbers below the panel indicate the ratio of *RAG1*:*18S*.

by the *HKK2* gene, is involved in glucose repression (RANDEZ-GIL *et al.* 1998). In fact, nuclear localization of Hxk2 appears to be glucose regulated, and it interacts *in vivo* with Mig1, the transcriptional repressor of many glucose-repressed genes (AHUATZI *et al.* 2004). In addition, in *K. lactis* and *S. cerevisiae*, hexokinase is also required for full glucose induction of *RAG1* and *HXT* gene expression (PRIOR *et al.* 1993; ÖZCAN and JOHNSTON 1999). Several pieces of evidence suggest that enolase is a multifunctional protein playing a crucial role in transcriptional and physiological processes (reviewed in PANCHOLI 2001). For example, enolase has been identified as a heat-shock protein in *S. cerevisiae* (HIDA and YAHARA 1985) and has been found to have transcriptional regulatory functions in larger eukaryotes. In the latter case, these functions occur through binding of enolase to certain gene promoters: *c-myc* (FEO *et al.* 2000; SUBRAMANIAN and MILLER 2000) or *STZ/ZAT10*, encoding a transcriptional repressor in Arabidopsis (LEE *et al.* 2002).

The finding that the expression of *RAG1* as well as several genes encoding glycolytic enzymes is affected in the $\Delta Kleno$ mutant suggests that enolase could play a regulatory role in *K. lactis* in addition to its catalytic activity. However, we have not yet demonstrated a direct role for enolase in the transcriptional regulation of *RAG1*. In addition to enolase it was already known that induction of *RAG1* expression by glucose is dependent on hexokinase activity (PRIOR *et al.* 1993). Since the loss of any one of the glycolytic steps tested, except that of phosphoglucose isomerase, severely reduces *RAG1* transcription, we believe that glucose metabolism generates a signal that induces *RAG1* expression. We cannot, however, exclude the possibility that enolase has a general regulatory function on other genes.

The defect in *RAG1* transcription in mutants blocked in glycolysis suggests the existence of regulatory mechanisms that prevent expression of genes encoding glucose transporters if a functional glycolytic pathway cannot be maintained. MILEOWSKI *et al.* (2001) previously showed that the absence of glucose transporters (hence glucose uptake) impaired the induction of *KHT1/RAG1* expression by high levels of glucose. Altogether, these data suggest that an intracellular glucose-sensing mechanism relying on glucose metabolism through glycolysis may ensure optimal glucose uptake by activating expression of the gene encoding the low-affinity glucose transporter. This intracellular pathway presumably collaborates with the extracellular glucose-sensing mechanism operating through the Rag4 glucose sensor in the cell membrane (BETINA *et al.* 2001). In *E. coli*, the expression of *ptsG*, encoding the major glucose transporter IIC^{Glc}, also requires glycolytic flux (KIMATA *et al.* 2001). Whatever the mechanism, regulation of glucose uptake by glycolytic flux seems to have been conserved from bacteria to yeasts.

We have a few clues to the mechanism by which glycolytic flux regulates expression of *RAG1* and glycolytic genes. The inactivation of *KIENO* leads to a severe reduction in expression of the regulatory genes *KIGCR1*, *KIGCR2*, *SCK1*, and *RAG4* (Figure 6). Thus, the effects of enolase on expression of genes encoding glycolytic enzymes and glucose transporters may be indirect. *KIGCR1*, *KIGCR2*, and *SCK1* are required for the full expression of glycolytic genes (LEMAIRE *et al.* 2002; NEIL *et al.* 2004). The KIGcr1/KIGcr2 complex directly regulates glycolytic gene expression through binding of KIGcr1 to glycolytic gene promoters (NEIL *et al.* 2004). An interaction between Sck1 and *K. lactis* glycolytic promoters is also probable since its *S. cerevisiae* ortholog




FIGURE 8.—Effect of the growth rate on *RAG1* transcription. (A) Inhibition of *K. lactis* growth by antimycin A, KCN, or G418. The MW270-7B (*KIENO*) strain was first grown in 50 ml YP medium containing 2% glycerol to an OD₆₀₀ of 2. Cells were harvested, washed, and resuspended in sterile cold water and aliquots (~15 OD₆₀₀) were diluted in 50 ml YPG (2% glucose) or YPG containing 5 μ M antimycin A, 5 mM KCN, or 100 μ g/ μ l G418. Cultures were kept agitated at 28° during 10 hr and OD₆₀₀ was checked at 0, 3, 6, and 10 hr. (B) Northern blot analysis of *RAG1*, *KIAC1*, and *18S* transcript levels. Total RNA was extracted from the YPG culture at *t* = 0 (shift on glucose) and from the four cultures at *t* = 10 hr. Electrophoresis conditions and quantification were done as described in the Figure 6 legend. The probes used to detect *RAG1*, *KIAC1*, and *18S* transcripts are described in MATERIALS AND METHODS. *KIAC1* and *18S* mRNA was used as internal standards and numbers below the corresponding panels indicate the ratio of *RAG1*:*KIAC1* or *RAG1*:*18S*.

(Sgc1) binds to the *ENO1* promoter (SATO *et al.* 1999). Hence, the transcriptional defect of *RAG5*/*RAG6* genes in the *Kleno* mutant is almost certainly the consequence of the low level of expression of *KIGCR1*, *KIGCR2*, and *SCK1*. Interestingly, these transcription factors are also required for the full glucose induction of *RAG1* expression (LEMAIRE *et al.* 2002; NEIL *et al.* 2004). At least in the case of *KIGcr1*/*KIGcr2*, this effect seems indirect since *KIGcr1* does not bind the *RAG1* promoter *in vitro* (NEIL *et al.* 2004) and thus may be a consequence of

the reduced glycolytic flux in the *KIGcr1*/*KIGcr2* mutants. Although we cannot exclude a cumulative effect of the reduced expression of *KIGCR1*, *KIGCR2*, *SCK1*, and *RAG4* on *RAG1* expression, we favor the hypothesis that a product of glycolysis may control the activity of an unidentified regulator. Such metabolic controls have been described already in *S. cerevisiae*. For instance, the Mcm1 transcriptional regulator is regulated post-transcriptionally by the glycolytic flux (CHEN and TYE 1995). Moreover, the efficient transcription of yeast AMP biosynthetic genes requires interaction between the transcription factors Bas1p and Bas2p, and this interaction is promoted in the presence of a metabolic intermediate (SAICAR) of this biosynthetic process (REBORA *et al.* 2001).

On the basis of the present findings, we propose that the regulation of *RAG1* expression by glucose involves two pathways: (i) a pathway involving the glucose sensor Rag4 that responds to extracellular glucose availability (BETINA *et al.* 2001) and (ii) a pathway responding to an intracellular signal generated by glycolysis. Together, these pathways can be considered to be an autoregulatory device for the fermentative utilization of sugars in yeast. The *K. lactis* system with its nonredundancy of the genes of glucose metabolism, the clear Rag[−] phenotype associated with glycolytic mutations, and its metabolic properties appears to be a suitable tool to study intracellular glucose sensing in yeast.

We thank Melanie Annat for cloning of the *KIENO* gene and Hiroshi Fukuhara for critical reading of the manuscript.

LITERATURE CITED

- AHUATZI, D., P. HERRERO, T. DE LA CERA and F. MORENO, 2004 The glucose-regulated nuclear localization of Hexokinase 2 in *Saccharomyces cerevisiae* is Mig-dependent. *J. Biol. Chem.* **279**: 14440–14446.
- BETINA, S., P. GOFFRINI, I. FERRERO and M. WÉSOLOWSKI-LOUVEL, 2001 *RAG1* gene encodes a glucose sensor in *Kluyveromyces fragilis*. *Genetics* **158**: 541–548.
- BIANCHI, M. M., L. TIZZANI, M. DESTREUILLE, L. FRONTALI and M. WÉSOLOWSKI-LOUVEL, 1996 The “petite-negative” yeast *Kluyveromyces fragilis* has a single pyruvate decarboxylase gene. *Mol. Microbiol.* **19**: 22–36.
- BILLARD, P., S. MENART, J. BLAISONNEAU, M. BOLOTIN-FUKUHARA, H. FUKUHARA *et al.*, 1996 Glucose uptake in *Kluyveromyces fragilis*: role of the *HGT1* gene in glucose transport. *J. Bacteriol.* **178**: 5860–5866.
- BISSON, L. F., and D. G. FRAENKEL, 1983 Involvement of kinases in glucose and fructose uptake by *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. USA* **80**: 1730–1734.
- BISSON, L. F., D. M. COONS, A. L. KRUEKEBERG and D. A. LEWIS, 1993 Yeast sugar transporters. *Crit. Rev. Biochem. Mol. Biol.* **28**: 259–308.
- BLAISONNEAU, J., H. FUKUHARA and M. WÉSOLOWSKI-LOUVEL, 1997 The *Kluyveromyces fragilis* equivalent of casein kinase I is required for the transcription of the gene encoding the low-affinity glucose permease. *Mol. Gen. Genet.* **253**: 469–477.
- BOEKE, J. D., F. LACROUTE and G. R. FINK, 1984 A positive selection for mutants lacking orotidine-5'-phosphate decarboxylase activity in yeast 5-fluoro-orotic acid resistance. *Mol. Gen. Genet.* **197**: 345–346.
- CHEN, T., T. HIROKO, A. CHAUDHURI, F. INOSE, M. LORD *et al.*, 2000

- Multigenerational cortical inheritance of the Rax2 protein in orienting polarity and division in yeast. *Science* **290**: 1975–1978.
- CHEN, X. J., 1995 Low- and high-copy-number shuttle vectors for replication in the budding yeast *Kluyveromyces fragilis*. *Gene* **172**: 131–136.
- CHEN, X. J., M. WESOLOWSKI-LOUVEL and H. FUKUHARA, 1992 Glucose transport in the yeast *Kluyveromyces fragilis*. II. Transcriptional regulation of the glucose transporter gene *RAG1*. *Mol. Gen. Genet.* **33**: 97–105.
- CHEN, Y., and B. TYE, 1995 The yeast Mcm1 protein is regulated posttranscriptionally by the flux of glycolysis. *Mol. Cell. Biol.* **15**: 4631–4639.
- CLIFTON, D., S. B. WEINSTOCK and D. C. FRAENKEL, 1978 Glycolysis mutants in *Saccharomyces cerevisiae*. *Genetics* **88**: 1–11.
- COHEN, R., J. P. HOLLAND, T. YOKOI and M. J. HOLLAND, 1986 Identification of a regulatory region that mediates glucose-independent induction of the yeast enolase gene *ENO2*. *Mol. Cell. Biol.* **6**: 2287–2297.
- COHEN, R., T. YOKOI, J. P. HOLLAND, A. E. PEPPER and M. J. HOLLAND, 1987 Transcription of the constitutively expressed yeast enolase gene *ENO1* is mediated by positive and negative cis-acting regulatory sequences. *Mol. Cell. Biol.* **7**: 2753–2761.
- DOHMEN, R. J., A. W. M. STRASSER, C. B. HÖNER and C. P. HOLLENBERG, 1991 An efficient transformation procedure enabling long-term storage of competent cells of various yeast genera. *Yeast* **7**: 691–692.
- ENTIAN, K. D., and K. L. FRÖHLICH, 1984 *Saccharomyces cerevisiae* mutants provide evidence of hexokinase PII as a bifunctional enzyme with catalytic and regulatory domains for triggering carbon catabolite repression. *J. Bacteriol.* **158**: 29–35.
- FEU, S., D. ARCURI, E. PIZZINI, R. PASSANTINO and A. GIALONGO, 2000 *ENO1* gene product binds to the *c-myc* promoter and acts as a transcriptional repressor: relationship with Myc promoter-binding protein (MBP-1). *FEBS Lett.* **473**: 47–52.
- GOFFRINI, P., A. A. ALGERI, C. DONNINI, M. WESOLOWSKI-LOUVEL and I. FERRERO, 1989 *RAG1* and *RAG2*: nuclear genes involved in the dependence/independence on mitochondrial respiratory function for the growth on sugars. *Yeast* **5**: 99–106.
- GOFFRINI, P., M. WESOLOWSKI-LOUVEL and I. FERRERO, 1991 A phosphoglucose isomerase gene is involved in the Rag phenotype of the yeast *Kluyveromyces fragilis*. *Mol. Gen. Genet.* **228**: 401–409.
- GONZALEZ SIRO, M. I., M. A. FREIRE PICOS and M. E. CERDAS, 1996 Reoxidation of the NADPH produced by the pentose phosphate pathway is necessary for the utilization of glucose by *Kluyveromyces fragilis* *rag2* mutants. *FEBS Lett.* **387**: 7–10.
- HAW, R., A. D. YARRAGUDI and H. UEMURA, 2001 Isolation of *GCRI*, a major transcription factor of glycolytic genes in *Saccharomyces cerevisiae*, from *Kluyveromyces fragilis*. *Yeast* **18**: 729–735.
- HOLLAND, M. J., J. P. HOLLAND, G. P. THILL and K. A. JACKSON, 1981 The primary structures of two yeast enolase genes. *J. Biol. Chem.* **256**: 1385–1395.
- IDA, H., and I. YAHARA, 1985 Yeast heat-shock protein of M_r 48,000 is an isoprotein of enolase. *Nature* **315**: 688–690.
- JACOBY, J., C. P. HOLLENBERG and J. HEINISCH, 1993 Transketolase mutants in the yeast *Kluyveromyces fragilis* provide evidence that glucose can be metabolized through the pentose phosphate pathway. *Mol. Microbiol.* **10**: 867–876.
- KIMATA, K., Y. TANAKA, T. INADA and H. AIBA, 2001 Expression of the glucose transporter gene, *ptsG*, is regulated at the mRNA degradation step in response to glycolytic flux in *Escherichia coli*. *EMBO J.* **20**: 3587–3595.
- LEE, H., Y. GUO, M. OHTA, L. XIONG, B. STEVENSON *et al.*, 2002 *LOS2*, a genetic locus required for cold-responsive gene transcription encodes a bi-functional enolase. *EMBO J.* **21**: 2692–2702.
- LEMAIRE, M., A. GUYON, S. BETINA and M. WESOLOWSKI-LOUVEL, 2002 Regulation of glycolysis by casein kinase I (Rag8p) in *Kluyveromyces fragilis* involves a DNA-binding protein, Sck1p, a homologue of Sck1p of *Saccharomyces cerevisiae*. *Curr. Genet.* **40**: 355–364.
- MCALISTER, L., and M. J. HOLLAND, 1982 Targeted deletion of a yeast enolase structural gene. *J. Biol. Chem.* **257**: 7181–7188.
- MILKOWSKI, C., S. KRAMPE, J. WEIRICH, V. HASSE, E. BOHLES *et al.*, 2001 Feedback regulation of glucose transporter gene transcription in *Kluyveromyces fragilis* by glucose uptake. *J. Bacteriol.* **183**: 5223–5229.
- NEIL, H., M. LEMAIRES and M. WESOLOWSKI-LOUVEL, 2004 Regulation of glycolysis in *Kluyveromyces fragilis*: role of *KIGCR1* and *KIGCR2* in glucose uptake and catabolism. *Curr. Genet.* **45**: 129–139.
- NIEDENTHAL, R., L. NILES, U. GULDENER, S. KLEIN, M. JOHNSTON *et al.*, 1999 Systematic analysis of *S. cerevisiae* chromosome VIII genes. *Yeast* **16**: 1775–1796.
- ÖZCAN, S., and M. JOHNSTON, 1999 Function and regulation of yeast hexose transporters. *Microbiol. Mol. Biol. Rev.* **63**: 554–569.
- PANCHOLI, V., 2001 Multifunctional α -enolase: its role in diseases. *Cell. Mol. Life Sci.* **58**: 902–920.
- PRIOR, C., P. MAMESSIER, H. FUKUHARA, X. J. CHEN and M. WESOLOWSKI-LOUVEL, 1993 The hexokinase gene is required for the transcriptional regulation of the glucose transporter gene *RAG1* in *Kluyveromyces fragilis*. *Mol. Cell. Biol.* **13**: 3882–3889.
- RANDEZ-GIL, F., P. SANZ, K. ENTIAN and J. A. PRIETO, 1998 Carbon source-dependent phosphorylation of Hexokinase PII and its role in the glucose-signaling response in yeast. *Mol. Cell. Biol.* **18**: 2940–2948.
- REBORA, K., C. DESMOUCELLES, F. BORNE, B. PINSON and B. DAIGNAN-FORNIER, 2001 Yeast AMP pathway genes respond to adenine through regulated synthesis of a metabolic intermediate. *Mol. Cell. Biol.* **23**: 7901–7912.
- SAFO, T., M. C. LOPEZ, S. SUGIYAMA, Y. JIGAMI, H. BAKER *et al.*, 1999 The E-box DNA binding protein Sgc1p suppresses the *ger2* mutation, which is involved in transcriptional activation of glycolytic genes in *Saccharomyces cerevisiae*. *FEBS Lett.* **463**: 307–311.
- SIBORSKY, R. S., and P. HIETER, 1989 A system of shuttle vectors and yeast host strains design for efficient manipulation of DNA in *Saccharomyces cerevisiae*. *Genetics* **122**: 19–27.
- SUBRAMANIAN, A., and D. M. MILLER, 2000 Structural analysis of α -enolase. *J. Biol. Chem.* **275**: 5958–5965.
- SUN, J. K., L. L. POULSEN, D. M. ZIEGLER and J. D. ROBERTUS, 1999 Yeast flavin-containing monooxygenase generates oxidizing equivalents that control protein folding in the endoplasmic reticulum. *Proc. Natl. Acad. Sci. USA* **96**: 2687–2691.
- WACH, A., 1996 PCR-synthesis of marker cassettes with long flanking homology regions for gene disruptions in *S. cerevisiae*. *Yeast* **12**: 259–265.
- WESOLOWSKI, M., A. A. ALGERI, P. GOFFRINI and H. FUKUHARA, 1982 Killer DNA plasmids of the yeast *Kluyveromyces fragilis*. I. Mutations affecting the killer phenotype. *Curr. Genet.* **150**: 137–140.
- WESOLOWSKI-LOUVEL, M., P. GOFFRINI, I. FERRERO and H. FUKUHARA, 1992a Glucose transport in the yeast *Kluyveromyces fragilis*. I. Properties of an inducible low-affinity glucose transporter gene. *Mol. Gen. Genet.* **33**: 89–96.
- WESOLOWSKI-LOUVEL, M., C. PRIOR, D. BORNECQUE and H. FUKUHARA, 1992b *Rag*⁺ mutations involved in glucose metabolism in yeast: isolation and genetic characterization. *Yeast* **8**: 711–719.

Communicating editor: M. JOHNSTON

Impact of Mitochondrial Function on Yeast Susceptibility to Antifungal Compounds

M. ŠARINOVÁ, E. TICHÁ, M. OBERNAUEROVÁ, Y. GBELSKÁ*

Department of Microbiology and Virology, Faculty of Science, Comenius University, 842 15 Bratislava, Slovakia

fax +421 265 429 064

e-mail gbelaska@fns.uniba.sk

Received 12 September 2006

Revised version 19 January 2007

ABSTRACT. *Saccharomyces cerevisiae pel1* and *crd1* mutants deficient in the biosynthesis of mitochondrial phosphatidylglycerol (PG) and cardiolipin (CL) as well as *Kluyveromyces lactis* mutants impaired in the respiratory chain function (RCF) containing dysfunctional mitochondria show altered sensitivity to metabolic inhibitors. The *S. cerevisiae pel1* mutant displayed increased sensitivity to cycloheximide, chloramphenicol, oligomycin and the cell-wall perturbing agents caffeine, caspofungin and hygromycin. On the other hand, the *pel1* mutant was less sensitive to fluconazole, similarly as the *K. lactis* mutants impaired in the function of mitochondrial cytochromes. Mitochondrial dysfunction resulting either from the absence of PG and CL or impairment of the RCF presumably renders the cells more resistant to fluconazole. The increased tolerance of *K. lactis* respiratory chain mutants to amphotericin B, caffeine and hygromycin is probably related to a modification of the cell wall.

Abbreviations

CL	cardiolipin		PG	phosphatidylglycerol	
mtDNA	mitochondrial DNA		PGPS	phosphatidylglycerophosphate synthase	
MIC	minimum inhibitory concentration		RCF	respiratory chain function	
Amb	amphotericin B	Cyh	cycloheximide	Ket	ketoconazole
Bif	bifonazole	Eco	econazole	Mic	miconazole
Caf	caffeine	Flu	fluconazole	Muc	mucidin
Chl	chloramphenicol	Hyg	hygromycin B	4-NQO	4-nitroquinoline <i>N</i> -oxide
Clo	clotrimazole	Itr	itraconazole	Oli	oligomycin
Cpf	caspofungin				

Mitochondria play a fundamental role in eukaryotic cell physiology producing cellular energy and other essential metabolites. By integrating numerous death signals mitochondria are also involved in the control of apoptosis. The biogenesis of functional mitochondria depends on the coordinated expression of two genomes, nuclear and mitochondrial, a cross-talk essential for cell life and death. While mitochondria contain their own DNA (mtDNA) encoding a handful of proteins, the vast majority of mitochondrial proteins are synthesized on cytosolic ribosomes and imported post-translationally into the organelle.

Mitochondrial genome integrity is essential for the viability of most yeast species. Loss of mtDNA or mitochondrial protein synthesis is lethal for *petite*-negative yeast species, e.g., *Kluyveromyces lactis*. The *K. lactis* cells are not able to tolerate the absence of both electron-proton transport pumping and ATP synthesis components of oxidative phosphorylation (Clark-Walker and Chen 2001). However, such a profound change can be tolerated by *petite*-positive *Saccharomyces cerevisiae*, readily forming respiratory-deficient *petite* mutants that lost mtDNA after treatment with DNA targeting drugs (Bulder 1964). On the other hand, mutations in several nuclear genes transform *S. cerevisiae* into a *petite*-negative species which no longer tolerates the cytoplasmic “*petite*” state: *opy1/aac2*, deficient in ATP/ADP translocase (Kolarov *et al.* 1990), *pel1/pgs1* deficient in PGPS (Janitor *et al.* 1996; Chang *et al.* 1998). The *pel1* (*petite* lethal) mutation, initially described as a nuclear mutation displaying synthetic lethal phenotype with mitochondrial *rho* mutations in *S. cerevisiae* (Šubík 1974), was found to be associated with CL deficiency (Janitor *et al.* 1996). It has been demonstrated that *PEL1* (renamed *PGS1*) encodes PGPS, the 1st enzyme in the CL biosynthesis pathway (Chang *et al.* 1998). The Pel1 and/or Pgs1p is localized to mitochondria, where both PG and CL are synthesized (Džugasová *et al.* 1998; Daun *et al.* 1998; Schlame *et al.* 2000). Disruption of *PEL1* (*PGS1*) results in complete loss of both PG and CL. CL (a phospholipid found predominantly in the inner mitochon-

*Corresponding author.

drial membrane) is essential for the biogenesis and proper RCF (Jiang *et al.* 2000; Vreken *et al.* 2000; Pfeiffer *et al.* 2003), ATP synthase (Koshkin and Greenberg 2002) and of ATP/ADP translocase (Hoffmann *et al.* 1994). PG can substitute for some essential functions of CL, both PG and CL being required for maintaining mtDNA (Džugasová and Šubík 2005; Zhong *et al.* 2005).

Due to the connection between the functional state of mitochondria and the activation of pleiotropic drug resistance network in *S. cerevisiae* (Hallstrom and Moye-Rowley 2000; Devaux *et al.* 2002), our aim was to analyze the possible influence of mitochondrial dysfunction resulting from altered mitochondrial membrane lipid composition in *petite*-positive *S. cerevisiae* as well as impaired RCF in *petite*-negative *K. lactis* on the cell's sensitivity to metabolic inhibitors and antifungal compounds.

MATERIAL AND METHODS

Media, culture conditions and strains. For the yeast strains used see Table I. Cells were grown at 30 °C in YPD complex medium with glucose (in %: glucose 2, bacteriological peptone 2, yeast extract 1), in YPG complex glycerol medium (in %: glycerol 2, bacteriological peptone 2, yeast extract 1) or YNB minimal medium (Difco) containing 0.67 % Yeast Nitrogen Base without amino acids, 2 % glucose, and supplemented with the auxotrophic requirements (40 mg/L). Solid media were prepared in 2 % Difco agar (all percentages are *M/V*). Respiration-deficient *petite* mutants from *S. cerevisiae* *crd1* strain were prepared by ethidium bromide (25 µmol/L) mutagenesis (Slonimski *et al.* 1968).

Table I. Yeast strains used

Strain	Genotype	Reference
<i>Saccharomyces cerevisiae</i>		
DW4-2A/YEp352/ <i>PEL1</i>	MATa <i>ade2-1 ura3-1 trp1-1 his3-11 pel1</i> /YEp352/ <i>PEL1</i>	Janitor and Šubík 1993
DW4-2A/YEp352 (<i>pel1</i>)	MATa <i>ade2-1 ura3-1 trp1-1 his3-11 pel1</i> /YEp352	this study
FY1679 (<i>CRD1</i>)	MATa <i>ura3-52 trp1Δ63 his3 Δ200 CRD1</i>	Tuller <i>et al.</i> 1998
FYC/5 (<i>Δcrd1</i>)	MATa <i>ura3-52 leu2 trp1Δ63 his3 Δ200 crd1::kanMX4</i>	ditto
FYC/5 (<i>Δcrd1/rho⁻</i>)	MATa <i>ura3-52 trp1Δ63 his3 D200 crd1::kanMX4/rho⁰</i>	this study
<i>Kluyveromyces fragilis</i>		
JBD100	a <i>trp1 lac4-1 ura3-100 Rag⁺</i>	Heus <i>et al.</i> 1990
JBD100/M1	a <i>trp1 lac4-1 ura3-100 gyl1 Rag⁺</i>	Gbelská <i>et al.</i> 1996
JBD100/M5	a <i>trp1 lac4-1 ura3-100 gyl2-2 Rag⁺</i>	ditto
JBD100/M7	a <i>trp1 lac4-1 ura3-100 gyl3-2 Rag⁺</i>	ditto
PM6-7A	a <i>ura41 ade2 Rag⁺</i>	Chen <i>et al.</i> 1992
PM6-7A/ <i>Δcox18</i>	a <i>ura41 ade2K1 cox18::URA3</i>	Hikkel <i>et al.</i> 1997

Drug susceptibility was tested qualitatively by spot assay. The strains were grown overnight at 30 °C in liquid YNB medium, cells were diluted to a concentration of 10⁷/nL (*i.e.* 10⁷ cells per nL) in sterile water and 10 nL of the cell suspension and 10-fold serial dilutions of cells were spotted onto complex and/or minimal plates with glucose or glycerol supplemented with various concentrations of the drugs tested, followed by incubation (3–5 d, 30 °C). Susceptibility to metabolic inhibitors, *i.e.* Cyh (*Sigma*) dissolved in ethanol, 4-NQO (*Sigma*), dissolved in acetone, Hyg (*Boehringer Mannheim*), Caf (*Fluka*), dissolved in Me₂SO, and antifungal compounds Cpf, Flu (*Pfizer Amboise*, France), dissolved in sterile water, Ket (*Janssen Pharmaceuticals*, Belgium), Mic (*Sigma*), Clo (*Sigma*), Bif (*Sigma*) dissolved in Me₂SO, was tested on YNB medium with glucose excepting Oli (*Sigma*) and Muc (*Sigma*) dissolved in ethanol, and Cfm (*Sigma*) dissolved in Me₂SO, that were tested on YPG. Sensitivity was also assessed using zone-inhibition assays. Approximately 10⁷ stationary-phase cells were plated onto minimal glucose agar. Filter discs (Ø 6 mm) soaked with an appropriate amount of antifungal compounds were placed on the plates, which were then incubated at 30 °C for 3–5 d before determining the diameter of the zone of growth inhibition.

Plasmid and transformation procedure. Cells of the strain DW4-2A were transformed using the empty high-copy episomal plasmid YEp352 (2 μ m, *URA3*, *amp^R*) or plasmid YEp352/*PEL1* containing the standard allele of the *PEL1* gene (Džugasová *et al.* 1998) using the lithium acetate procedure (Sambrook *et al.* 1989).

RESULTS AND DISCUSSION

S. cerevisiae mutants deficient in PG and CL are sensitive to metabolic inhibitors. Sensitivity to a broad variety of metabolic inhibitors correlates with mitochondrial function (Zhang and Moye-Rowley 2001; Devaux *et al.* 2002). CL and PG (anionic phospholipids) are synthesized inside the mitochondria, essential for the proper function of the organelle. Therefore the sensitivity of *pel1* mutant (deficient in PG and CL), *crd1* mutant (deficient in CL) and *crd1/rho* mutant (deficient in CL and mtDNA) to several metabolic inhibitors (Cyh, Cln, 4-NQO, Oli, Muc) was compared with that of the corresponding wild-type strains *pel1*/YEp352 – *PEL1* and *CRD1*, respectively. The simultaneous absence of PG and CL in the *pel1* mutant renders the cells more sensitive to Cyh, Oli and Cln in comparison with the corresponding wild-type strain (Table II). The MIC of Muc was similar for both mutants (*pel1*, *crd1*) and the corresponding wild-type

Table II. Sensitivity of *S. cerevisiae* strains to metabolic inhibitors (MIC, μ g/mL)

Strain	Cyh ^a	4-NQO ^b	Oli ^c	Cln ^d	Muc ^e
DW4-2A/YEp352/ <i>PEL1</i>	0.15	0.3	>1	>14	<0.025
DW4-2A/YEp352 (<i>pel1</i>)	0.05	0.2	0.6	12	<0.025
FY1679 (<i>CRD1</i>)	0.1	0.2	>0.9	14	0.05
FYC5 (Δ <i>crd1</i>)	0.1	0.1	0.9	12	<0.025
FYC5 (Δ <i>crd1/rho</i>)	0.1	0.2	–	–	–

^aCycloheximide 0, 0.05, 0.1, 0.15, 0.2, 0.25, 0.3–0.7

^dChloramphenicol 5–14

^b4-Nitroquinoline *N*-oxide 0.1, 0.2, 0.3, 0.4–0.7

^eMucidin 0.025, 0.05, 0.075, 0.1

^cOligomycin 0.1–0.9

strains. On the other hand, the deficiency of CL alone in the *crd1* mutant or the simultaneous absence of CL and mtDNA in the double *crd1/rho* mutant did not influence the sensitivity of cells to the metabolic inhibitors. The observed effect in the *pel1* mutant may be associated with an increased drug efflux induced by dysfunctional mitochondria. The loss of mitochondrial function led to the activation of the *PDR3* gene-mediated overexpression of *PDR5* encoding a multidrug resistance efflux pump (Zhang and Moye-Rowley 2001; Devaux *et al.* 2002).

However, the presence of PG and CL is essential for physiology (Janitor and Šubík 1993; Chang *et al.* 1998; Džugasová *et al.* 1998; Gohil *et al.* 2004) and biogenesis of mitochondria (Jaing *et al.* 2000; Schlame *et al.* 2000; Ostrander *et al.* 2001; Koshkin and Greenberg 2002) and their protective role in the eukaryotic cell response to oxidative stress was also proposed (Mileykovskaya *et al.* 2005).

Mitochondrial dysfunction affects the susceptibility of petite-positive and petite-negative yeasts to antifungal compounds. The deficiency in PG and CL rendered the *S. cerevisiae pel1* mutant cells more sensitive to Ket, Mic, Bif and Amb (Table III) in comparison with the isogenic wild-type strain. Moderate increase in sensitivity to azole antifungals was also seen in the *S. cerevisiae* Δ *crd1* mutant deficient in CL only in comparison with its parental wild-type strain (*CRD1*). However, the loss of mtDNA in the Δ *crd1* mutant caused an increased sensitivity to the antifungal compounds. The *pel1* mutant deficient in PG and CL was less sensitive to Flu than its isogenic transformant containing the *PEL1* gene on a multicopy plasmid (Table III, Fig. 1). The presence of PG in the *S. cerevisiae* Δ *crd1* mutant could substitute the essential functions of CL in mitochondria and the cells behave more or less like wild-type ones. The absence of both PG and CL in the *pel1* mutant leads to mitochondrial dysfunction that is associated with the observed Flu resistance.

The sensitivity of *K. lactis* mitochondrial mutants to antifungal compounds depends on RCF. The *K. lactis* nuclear mutants deficient in cytochrome *c*, cytochrome *c*₁ and cytochrome oxidase were less sensitive to Flu, Mic, Clo, Ket, Bif, and Eco in comparison with the corresponding wild-type strain (Fig. 2). The results of disk diffusion assays correlated with those observed in the qualitative testing of cell sensitivity by spotting serial dilutions of cells onto YPD or minimal agar plates containing different concentrations of anti-

Table III. Sensitivity of *S. cerevisiae* strains to antifungal compounds and metabolic inhibitors (MIC)

Strain	Flu ^a µg/mL	Ket ^b µg/mL	Mic ^c µg/mL	Clo ^d µg/mL	Bif ^e µg/mL	Amb ^f µg/mL	Hyg ^g mg/mL	Caf ^h mg/mL	Cpf ⁱ mg/mL
DW4-2A/YEp352/ <i>PEL1</i>	<14	7	0.1	0.6	5	7.5	0.1	>1	>0.75
DW4-2A/YEp352 (<i>pel1</i>)	>15	6	<0.1	0.5	2.5	5	0.1	<1	0.5
FY1679 (<i>CRD1</i>)	>15	6	0.1	0.6	>5	5	0.4	2	0.7
FYC5 (<i>Δcrd1</i>)	12	2	<0.1	0.6	>5	5	0.4	2	0.5
FYC5 (<i>Δcrd1 rho</i>)	12	<2	<0.1	0.4	2.5	2.5	0.2	<1	0.25

^aFluconazole 0, 10–15 µg/mL^dClotrimazole 0.1–0.8 µg/mL^gHygromycin 0.1–1 mg/mL^bKetoconazole 1–10 µg/mL^eBifonazole 2.5, 5, 7.5, 10 µg/mL^hCaffeine 0, 0.5, 1, 1.5, 2, 2.5, 3 mg/mL^cMiconazole 0.1–0.8 µg/mL^fAmphotericin B 0, 2.5, 5, 7.5, 10 µg/mLⁱCaspofungin 0, 0.1, 0.25, 0.5, 0.75 mg/mL

fungal compounds. All of the *K. lactis* mutants affected in the RCF were able to grow on plates containing Flu at 15 µg/mL in contrast with the isogenic wild-type strains (Fig. 1). Taken together, we found that the loss of mitochondrial function resulting from the absence of PG and CL in the *S. cerevisiae pel1* mutant or impaired RCF in *K. lactis* mutants is linked to the increased tolerance of yeast cells to Flu.




Fig. 1. Fluconazole sensitivity profiles of *S. cerevisiae* (A) and *K. lactis* (B) mutants. Wild type and mutant strains were grown in YNB medium for overnight at 30 °C, the A_{600} was normalized and 10 µL of 10-fold serial dilutions were spotted onto YNB plates with or without fluconazole, photographed after 5 d of growth at 30 °C.

Flu acts as ergosterol-depleting agent in *S. cerevisiae* and mitochondria could function as important physiological partners in the accumulation of toxic sterol intermediates in the presence of azoles (Kontoyianis 2000). Dysfunctional mitochondria could not assist in the accumulation of toxic sterol intermediates and the cells can tolerate the presence of Flu.

Mitochondrial functions are linked to cell-wall biogenesis. Zhong *et al.* (2005) indicated the role of mitochondrial lipids PG and CL in mitochondrial functions required for cell-wall biogenesis. Therefore the sensitivity of *S. cerevisiae* mutants deficient in PG and CL to Caf, Cpf and Hyg were assessed. The absence of PG and CL in *S. cerevisiae pel1* mutant renders the cells more sensitive to Caf and Cpf in comparison with the isogenic wild-type strain (Table III). The sensitivity of *S. cerevisiae Δcrd1* mutant and its corresponding wild-type strain to the inhibitory compounds used were similar. The presence of PG can apparently substitute for the CL functions in the *Δcrd1* mutant. The loss of mtDNA in the *Δcrd1* mutant caused even higher sensitivity of cells to Caf, Hyg and Cpf. The results correlate with the proposed role of PG and CL in functions that are essential for cell wall biogenesis. On the other hand, the *K. lactis* mutants affected in the RCF (deficient in cytochrome *c*, cytochrome *c*₁ or cytochrome oxidase) showed an increased tolerance to




Fig. 2. Antifungal sensitivity of wild-type *K. lactis* PM6-7A and PM6-7A/ Δ cox18 mutant cells; performed by the disc diffusion assay on YNB agar plates; Eco – econazole, Flu – fluconazole, Mic – miconazole, Cln – clotrimazole, Ket – ketoconazole.

Amb, Caf and Hyg (Table IV) suggesting changes in the cell wall structure as a result of mitochondrial dysfunction. The correlation between mutations in genes related to mitochondrial functioning and yeast cell wall characteristics was first mentioned in 1980 (Evans *et al.* 1980). Later on the connection between Amb

Table IV. Sensitivity of *K. lactis* strains to antifungal compounds and metabolic inhibitors (MIC)

Strain	Flu ^a μg/mL	Bif ^b μg/mL	Itc ^c μg/mL	Amb ^d μg/mL	Hyg ^e μg/mL	Caf ^f mg/mL
JBD100 (wt)	20	3	7.5	0.5	25	1
JBD100/M1	>50	>10	>50	>2	50	3
JBD100/M5	>50	>10	>50	>2	50	3
JBD100/M7	>50	>10	>50	>2	50	3
PM6-7A (wt)	20	4	15	1	50	2
PM6-7A/ Δ cox18	50	6	>15	>2	50	3

^aFluconazole 10, 20, 25, 30–50 μg/mL

^bBifonazole 0.5, 1, 1.5, 2–6, 10 μg/mL

^cItraconazole 2.5, 5, 7.5, 10, 15, 20, 25, 30, 50 μg/mL

^dAmphotericin B 0, 0.5, 1, 2 μg/mL

^eHygromycin 5, 10, 25, 50 μg/mL

^fCaffeine 0, 1–4 mg/mL

resistance and modification of the cell wall in *S. cerevisiae* (Lussier *et al.* 1997; Wauters *et al.* 2001; Zhang *et al.* 2003; Agarwal *et al.* 2003; Hapala *et al.* 2005) and also in *K. lactis* has been reported (Ramanandraibe *et al.* 1998). The results obtained using both *petite*-positive and *petite*-negative species show that the functional state of yeast mitochondria influences the surface characteristics of the cell as well as the susceptibility of cells to antifungal compounds.

The work was supported by grants of the Slovak Grant Agency VEGA (1/2338/05, 1/3246/06).

REFERENCES

- AGARWAL A.K., ROGERS P.D., BAERSON S.R., JACOB M.R., BARKER K.S., CLEARY J.D., WALKER L.A., NAGLE D.G., CLARK A.M.: Genome-wide expression profiling of the response to polyene, pyrimidine, azole and echinocandin antifungal agents in *Saccharomyces cerevisiae*. *J.Biol.Chem.* 278, 34998–35015 (2003).
- BULDER C.J.A.E.: Induction of *petite* mutations and inhibition of synthesis of respiratory enzymes in various yeasts. *Antonie van Leeuwenhoek J.Microbiol.* 30, 1–9 (1964).

- CHANG S.C., HAECOCK P.N., CLANCY C.J., DOWHAN W.: The *PEL1* gene (renamed *PGS1*) encodes the phosphatidylglycerophosphate synthase of *Saccharomyces cerevisiae*. *J.Biol.Chem.* 273, 9829–9839 (1998).
- CHEN X.J., WESOŁOWSKI-LOUVEL M., FUKUHARA H.: Glucose transport in the yeast *Kluyveromyces fragilis*. II. Transcriptional regulation of the glucose transporter gene *RAG1*. *Mol.Gen.Genet.* 233, 97–105 (1992).
- CLARK-WALKER G.D., CHEN X.J.: Dual mutations reveal interactions between components of oxidative phosphorylation in *Kluyveromyces fragilis*. *Genetics* 159, 929–938 (2001).
- DAUM G., LEES N.D., BARD M., DICKSON R.: Biochemistry, cell biology and molecular biology of lipids of *Saccharomyces cerevisiae*. *Yeast* 14, 1471–1510 (1998).
- DEVAUX F., CARVAJAL E., MOYE-ROWLEY W.S., JACQ C.: Genome-wide studies on the nuclear *PDR3*-controlled response to mitochondrial dysfunction in yeast. *FEBS Lett.* 515, 25–28 (2002).
- DŽUGASOVÁ V., OBERNAUEROVÁ M., HORVÁTHOVÁ K., VACHOVÁ M., ŽÁKOVÁ M., ŠUBÍK J.: Phosphatidylglycerophosphate synthase encoded by the *PEL1/PGS1* gene in *Saccharomyces cerevisiae* is localized in mitochondria and its expression is regulated by phospholipid precursors. *Curr.Genet.* 34, 297–302 (1998).
- DŽUGASOVÁ V., ŠUBÍK J.: Synthetic lethal interaction between the *pel1* and *opl* mutations in *Saccharomyces cerevisiae*. *Folia Microbiol.* 50, 293–299 (2005).
- EVANS I.H., DIALA E.S., EARL A., WILKIE D.: Mitochondrial control of cell surface characteristics in *Saccharomyces cerevisiae*. *Biochim.Biophys.Acta* 602, 201–206 (1980).
- GEELSKÁ Y., HORVÁTHOVÁ K., VAN DER AART Q.J.M., ZONNEVELD B.J.M., STEENSMAN H.Y., ŠUBÍK J.: Isolation and molecular analysis of the gene for cytochrome *c1* from *Kluyveromyces fragilis*. *Curr.Genet.* 30, 145–150 (1996).
- GOHIL V.M., HAYES P., MATSUYAMA S., SCHAGGER H., SCHLAME M., GREENBERG M.L.: Cardiolipin biosynthesis and mitochondrial respiratory chain function are interdependent. *J.Biol.Chem.* 279, 42612–42618 (2004).
- HALLSTROM T.C., MOYE-ROWLEY W.S.: Multiple signals from dysfunctional mitochondria activate the pleiotropic drug resistance pathway in *Saccharomyces cerevisiae*. *J.Biol.Chem.* 275, 37347–37356 (2000).
- HAPALA L., KLOBUČNIKOVÁ V., MAŽAŇOVÁ K., KOHÚT P.: Two mutants selectively resistant to polyenes reveal distinct mechanisms of antifungal activity by nystatin and amphotericin B. *Biochem.Soc.Trans.* 33, 1206–1209 (2005).
- HEUS J.J., ZONNEVELD B.J.M., STEENSMAN H.Y., VAN DEN BERG J.A.: Centromeric DNA of *Kluyveromyces fragilis*. *Curr.Genet.* 18, 517–522 (1990).
- HIKEL I., GEELSKÁ Y., VAN DER AART Q.J.M., LUBEC G., ŠUBÍK J.: Cloning and characterization of *KICOX18*, a gene required for activity of cytochrome oxidase in *Kluyveromyces fragilis*. *Curr.Genet.* 32, 267–272 (1997).
- HOFFMANN B., STOCK A., SCHLAME M., BEYER K., KLINGENBERG M.: The reconstituted ADP/ATP carrier activity has an absolute requirement for cardiolipin as shown in cysteine mutants. *J.Biol.Chem.* 269, 1940–1944 (1994).
- JANITOR M., ŠUBÍK J.: Molecular cloning of the *PEL1* gene of *Saccharomyces cerevisiae* that is essential for the viability of *petite* mutants. *Curr.Genet.* 24, 307–312 (1993).
- JANITOR M., OBERNAUEROVÁ M., KOHLWEIN S.D., ŠUBÍK J.: The *pel1* mutant of *Saccharomyces cerevisiae* is deficient in cardiolipin and does not survive the disruption of the *CHO1* gene encoding phosphatidylserine synthase. *FEMS Microbiol.Lett.* 140, 43–47 (1996).
- JIANO F., RYAN M.T., SCHLAME M., ZHAO M., GU Z., KLINGENBERG M., PFANNER N., GREENBERG M.L.: Absence of cardiolipin in the *crd1* null mutant results in decreased mitochondrial membrane potential and reduced mitochondrial function. *J.Biol.Chem.* 275, 22387–22394 (2000).
- KAUR R., CASTANO I., COORMACK B.P.: Functional genomic analysis of fluconazole susceptibility in the pathogenic yeast *Candida glabrata*: roles of calcium signalling and mitochondria. *Antimicrob.Agents Chemother.* 48, 1600–1613 (2004).
- KOLAROV J., KOLAROVÁ N., NELSON N.: A third ADP/ATP translocator gene in yeast. *J.Biol.Chem.* 265, 12711–12716 (1990).
- KONTOTIANNIS D.P.: Modulation of fluconazole sensitivity by the interaction of mitochondria and Erg3p in *Saccharomyces cerevisiae*. *J.Antimicrob.Chemother.* 46, 191–197 (2000).
- KOSHIN V., GREENBERG M.L.: Cardiolipin prevents rate-dependent uncoupling and provides osmotic stability in yeast mitochondria. *Biochem.J.* 364, 317–322 (2002).
- LUSSIER M., WHITE A.M., SHERATON J., DI PAOLO T., TREADWELL J., SOUTHARD S.B., HORENSTEIN C.I., CHEN-WEINER J., RAM A.F.J., KAPTEYN J.C., ROEMER T.W., VO D.H., BONDOC D.C., HALL J., ZHONG W.W., SDOU A.M., DAVIES J., KILS F.M., ROBERTS P.W., BUSSEY H.: Large scale identification of genes involved in cell surface biosynthesis and architecture in *Saccharomyces cerevisiae*. *Genetics* 147, 435–450 (1997).
- MILEYKOVSKAYA E., ZHANG M., DOWHAN W.: Cardiolipin in energy transducing membranes. *Biochemistry (Moscow)* 70, 154–158 (2005).
- OSTRANDER D.B., ZHANG M., MILEYKOVSKAYA E., RHO M., DOWHAN W.: Lack of mitochondrial anionic phospholipids causes an inhibition of translation of protein components of the electron transport chain. A yeast genetic model system for the study of anionic phospholipid function in mitochondria. *J.Biol.Chem.* 276, 25262–25272 (2001).
- PFEIFFER K., GOHIL V., STUART R.A., HUNTE C., BRANDT U., GREENBERG M.L.: Cardiolipin stabilizes respiratory chain supercomplexes. *J.Biol.Chem.* 278, 52873–52880 (2003).
- RAMANANDRAIBE E., YOUNSI M., COULON J., LOFFINET V., HAKKOU A., BONALY R.: Implication of cell wall constituents in the sensitivity of *Kluyveromyces fragilis* strains to amphotericin B. *Res.Microbiol.* 149, 109–118 (1998).
- SAMBROOK J., FRITSCH E.F., MANIATIS T.: *Molecular Cloning: a Laboratory Manual*, 2nd ed. Cold Spring Harbor Laboratory Press, New York 1989.
- SCHLAME M., RUA D., GREENBERG M.L.: The biosynthesis and functional role of cardiolipin. *Progr.Lipid Res.* 39, 257–288 (2000).
- SLONIMSKI P.P., PERRODIN G., CROFT J.H.: Ethidium bromide induced mutation of yeast mitochondria: complete transformation of cells into respiratory deficient non-chromosomal “petites”. *Biochem.Biophys.Res.Comm.* 30, 232–239 (1968).
- ŠUBÍK J.: A nuclear mutant of *S. cerevisiae* non-tolerating the cytoplasmic *petite* mutation. *FEBS Lett.* 42, 309–312 (1974).

- TULLER G., HRASTNIC C., ACHLEITNER G., SCHEFTHALER U., KLEIN F., DAUM G.: *YDL142c* encodes cardiolipin synthase (Clp1p) and is non-essential for aerobic growth of *Saccharomyces cerevisiae*. *FEBS Lett.* **421**, 15–18 (1998).
- VREKEN P., VALIANPOUR F., NUTMANS L.G., GRIVELL L.A., PLECKO B., WANDERS R.J., BARTH P.G.: Defective remodeling of cardiolipin and phosphatidylglycerol in Barth syndrome. *Biochem.Biophys.Res.Commun.* **20**, 378–382 (2000).
- WAUTERS T., ISEMENTANT D., VERACHTERT H.: Impact of mitochondrial activity on the cell wall composition and on the resistance to tannic acid in *Saccharomyces cerevisiae*. *J.Gen.Appl.Microbiol.* **47**, 21–26 (2001).
- ZHANG X., MOYE-ROWLEY W.S.: *Saccharomyces cerevisiae* multidrug resistance gene expression inversely correlates with the status of the F_0 component of the mitochondrial ATPase. *J.Biol.Chem.* **276**, 47844–47852 (2001).
- ZHANG M., SU X., MILEYKOVSKAYA E., AMOSCATO A., DOWHAN W.: Cardiolipin is not required to maintain mitochondrial DNA stability or cell viability for *Saccharomyces cerevisiae* grown at elevated temperatures. *J.Biol.Chem.* **278**, 35204–35210 (2003).
- ZHONG Q., GVOZDENOVIC-JEREMIC J., WEBSTER P., ZHOU J., GREENBERG M.L.: Loss of function of *KRE5* suppresses temperature sensitivity of mutants lacking mitochondrial anionic lipids. *Mol.Biol.Cell* **16**, 665–675 (2005).